



DIRECTIONS FOR USE

RESTRAINTS DO NOT apply by air.

DO NOT apply to waterlogged soil

DO NOT apply to fields where the slope exceeds 3%.

DO NOT apply if heavy rains or storms that are likely to cause surface runoff are forecast within 2 days of application

DO NOT irrigate to the point of runoff for at least 2 days after application.

DO NOT use as a pre-emergence application in sorghum during the wet season in the Northern irrigation areas of Western Australia

DO NOT apply more than 1.2 kg per crop.

Spray Drift Restraints:

DO NOT apply with spray droplets smaller than a COARSE spray droplet size category according to the "APVMA Compliance Instructions for Mandatory COARSE or Larger roplet Size Categories" located under this title in the GENERAL INSTRUCTIONS section of this label

FLAT SIZE: 332 MM WIDE X 448 MM DEEF

DO NOT apply when wind speed is less than 3 or more than 20 kilometres per hour as measured at the application site

DO NOT apply during surface temperature inversion conditions at the application site

Users of this product MUST make an accurate written record of the details of each spray application within 24 hours following application and KEEP this record for a minimum of 2 years. The spray application details that must be recorded are: 1) date with start and finish times of application; 2) location address and paddock/s sprayed; 3) full name of this product; 4) amount of product used per hectare and number of hectares applied to; 5) crop/situation and weed/pest; 6) wind speed and direction during application; 7) air temperature and relative humidity during application; 8) nozzle brand, type, spray angle nozzle capacity and spray system pressure measured during application; 9) name and address of person applying this product. (Additional record details may be required by the state or territory by the state or territory where this product is used.)

Mandatory No-Spray Zones for protection of the aquatic environment and non-target vegetation or animal habitat: **DO NOT** apply if there are aquatic or wetland areas including aquacultural ponds within 60 metres downwind from the application area. **DO NOT** apply if there are non-target vegetation or animal habitat within 30 metres downwind from the application area.

CROP	SITUATION	WEEDS	RATE	CRITICAL COMMENTS
Chickpeas, Faba beans, Field peas, Lupins. NSW, Vic, SA, WA, Qld only	Pre-Sowing	Suppression of: Wild Radish, Doublegee/ Spiny emex	0.86 to 1.2 kg/ha	Use the lower rate on light sandy soils (sandy loams to loamy sands) and the higher rate on heavier soils (loams, silt plus clay 40-60%). The soil should be free of excessive clods, trash and deep furrows. Sufficient rainfall (20-30 mm to wet the soil through the weed root zone is necessary within 2 to 3 weeks of application. In Pulse crops ensure the seed is covered with at least 3 cm and preferably 5 cm of soil and do not use rates higher than 0.86 kg/ha on soils with pH 8.0 and above as unacceptable crop damage may occur. NOTE: Some early crop phytotoxicity may be observed particularly on light soils. Heavy, intense rainfall following application may cause crop damage. At higher rates, avoid overlapping sprays and spraying-out corners.
	Post Sowing Pre- Emergence(PSPE). Apply to the soil within 2 days after sowing	Burr medic, Black bindweed, Corn gromwell (White ironweed, Sheepweed), Deadnettle, Indian hedge mustard, Long haired poppy, Mint weed, Ox tongue, Prickly lettuce, Shepherd's purse, Sow thistle, Toad rush, Turnip weed, Wild gooseberry, Wild turnip, Wireweed/hogweed. Suppression of: Annual ryegrass, Doublegee/Spiny emex, Phalaris, Wild oats, Wild radish	0.60 to 0.86 kg/ha	
Canola (TT or Triazine Tolerant varieties		Suppression of: Wild Radish, Doublegee/ Spiny emex	0.86 to 1.2 kg/ha	
only)	Post Sowing Pre-Emergence (PSPE). Apply to the soil within 2 days after sowing	Burr medic, Black bindweed, Corn gromwell (White ironweed, Sheepweed), Deadnettle, Indian hedge mustard, Long haired poppy, Mint weed, Ox tongue, Prickly lettuce, Shepherd's purse, Sow thistle, Toad rush, Turnip weed, Wild gooseberry, Wild turnip, Wireweed/hogweed. Suppression of: Annual ryegrass, Doublegee/ Spiny emex, Phalaris, Wild oats, Wild radish		
	Early Post-Emergence	Wild Radish, Deadnettle Suppression of: Annual ryegrass, Doublegee/Spiny emex, Phalaris, Wild oats	0.6 to 1.2 kg/ha	Apply when weeds are actively growing. Apply up to the 6 leaf stage of canola. Use higher rate on weeds 6-8 true leaves at time of application. Always add Hasten* at 500 mL/100 L water. DO NOT add any other herbicide adjuvant or wetter.

ciotning, and respirator and it rubber, wash with detergent and warm water. with water. Wash hands after use. After each day's use wash gloves, contaminated respirator with dust cartridge or canister. If product in eyes, wash it out immediately and a washable hat, elbow length chemical-resistant gloves and a half facepiece and loading and preparing spray, wear cotton overalls buttoned to the neck and wrist May irritate the eyes. Avoid contact with eyes. When opening the container, mixing SAFETY DIRECTIONS

Do not burn empty containers or product. marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulation. bury the empty packaging 500 mm below the surface in a disposal pit specifically recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility if an approved waste management facility is not available cap and return clean containers to recycler or designated collection point. If not TO THE Shigh ISTURY DO HOT AISHOSE OF UTUALIDING CHEMICS SON-SITE: IT LECYCHILD, TEPRACE sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings Store in the closed, original container in a dry, cool, well-ventilated area out of direct JASOASIG GNA 30AROTS

with susceptible crops or plants. DO NOT plant crops other than those recommended of this label for at least 6 months following the most recent application of up to 1.2water from heavy rain can be expected to run off to areas containing, or to be planted or on areas where their roots may extend or in locations where the chemical may be washed or moved into contact with their roots, DO NOT apply on sites where surface DO NOT apply or drain or flush equipment on or near desirable trees or other plants PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

water. DO NOT use in channels or drains. where turrows have been created in the soil for the purpose or holding or channelling 11-Canola: DO NOT use or apply this product post-emergence on raised beds or point of run-off, unless it can be retained on farm. product or used containers. After application of SPUTNIK, DO NOT irrigate crop to the Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with the PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre, Phone Australia

SDS

Additional information is listed in the safety data sheet (SDS). A safety data sheet for FARMALINX SPUTNIK Herbicide is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 9389 2455.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Farmalinx Pty Ltd or under abnormal conditions. Farmalinx Ptv Ltd accepts no liability for any loss or damage arising from incorrect



FARMALINX Pty Ltd ABN 95 134 353 245 Level 25, Suite 2506, Tower 2, 101 Grafton Street, Bondi Junction NSW 2022 T: 02 9389 2455 F: 02 9389 2844 Website: www.farmalinx.com

APVMA Approval No. 80864/118015

resistant gloves. Clothing must be laundered after each day's use. cotton overalls buttoned to the neck and wrist over normal clothing and chemical Do not allow entry into treated areas for 8 days. When prior entry is necessary, wear RE-ENTRY PERIOD

iabei spray uriti kestraint. specifies for the selected nozzle to deliver the droplet size category required in the DO NOT use a higher spray system pressure than the manufacturer ASAE S572 or BCPC. Choose a nozzle specified to provide the droplet size category required in the label Spray Driff Restraints.

a VERY CORRSE or an EXTREMELY COARSE droplet size category as referenced to USE ONLY nozzles that the nozzles' manufacturer has rated to deliver a CUARISE, Mandatory Instructions for Ground Applications

Instructions for Ground Application - for COARSE droplet size or larger a COARSE or larger droplet size category. ethods specified in these instructions to select a correct nozzle to deliver DRIY THE TOHOWING SPECIFIC METHODS FOR CHOOSING THE COFFECT HOZZIE. USE ONE OF under the spray operation conditions being used. The APVMA has approved Complying with the label requirement to use a specific droplet size category means using the correct nozzle that will deliver that droplet size category nozzle to comply with this requirement, the manufacturer must refer to at least one. manufacturers may refer to one or both to identify droplet size categories, but for a (newer name may also be shown as ASABE) or the BCPC guideline. Mozzle application. Spray droplet size categories are defined in the ASAE S572 Standard with the requirement of a COARSE or larger spray droplet size category for spray These instructions inform users of this chemical product how to lawfully comply mportant Information

APVMA Compliance Instructions for Mandatory COARSE or Larger Droplet Size spray quality according to ASAE S572 definition for standard nozzles. develop. Apply by ground boom only in a minimum of 50 L/ha. Use only COARSE bel none of dufing weather conditions when subject temperature inversions can DO NOT apply when wind speed is less than 3 and greater than 20 kilometres



GROUP

ABN 95 134 353 245 r.2, 101 Grafton Street, Ltd Tower 2022 FARMALINX Pty Li Level 25, Suite 2506, Tc Bondi Junction NSW 20

CROP SITUATION RATE CRITICAL COMMENTS WEEDS Lentils 0.86 to 1.2 kg/ha Lentils are more sensitive to SPUTNIK than other Pre-Sowing Suppression of: Wild radish, Doublegee/ Pulse crops. Do not use on light soils (ie, sandy loams to loamy sands, less than 40% clay). The crop must be sown at least 5 cm deep. Refer to APPLICATION section of the label. Use the lower rate on light weed infestations. Do not use rates higher than 0.86 kg/ha on soils with pH 8.0 and above as unacceptable crop damage may occur. The soil should be free of excessive clods, trash and deep furrows. Sufficient rainfall (20 to 30 mm) to wet the soil through the weed root zone is necessary within 2 to 3 weeks of application. NOTE: Some early crop phytotoxicity may be observed at the high rate. Heavy, intense rainfall following application may cause crop damage. Avoid over-lapping sprays and spraying-out Apply to soil pre-sowing or post-sowing. Refer to the APPLICATION section below. Sorghum Pre-Sowing AND Post-Bladder ketmia, Caltrop (yellow vine), 1.2 kg/ha Sowing Pre-Emergence Dwarf Amaranth, Milk thistle (sow thistle and Pigweed Always add Hasten at 500 ml /100 L water Apply Post -Emergence Boggabri weed, Noogoora burr, Sow 1.2 kg/ha up to the 5-7 leaf stage of sorghum. Refer to the thistle (milk thistle), Turnip weed, Wild fluroxypyr 200 g/L label for complete directions gooseberry. Always add Hasten at 500 mL/100 L water. Apply All weeds above plus-0.95 kg/ha Annual ground cherry, Apple of Peru, up to the 5-7 leaf stage of sorghum. Refer to the fluroxypyr 200 g/L label for complete directions Bathurst burr, Bladder ketmia, Caltrop 500-750 mL/ha of (vellow vine). Cow vine (peach vine). 200 a/L and rate selection. Dwarf amaranth, Giant (black) pigweed fluroxypyi Pigweed (red), Thornapple (Datura spp) 0.86 - 1.2 kg/ha African turnip weed, Burr medic, Black Apply to moist soil or when rain is imminent and Application to fallows | Pre-emergence bindweed, Corn gromwell (White iron prior to planting prior to germination of weeds. Use the higher weed, Sheep weed), Dead nettle, Indian mung beans, rate where the soil is covered by more than 50% hedge mustard. Flax leaf fleabane, Long sovbeans trash/stubble/weeds, for heavy weed pressure haired poppy, Mint weed, Ox tongue, or for longer weed control. If weeds are present, barley, oats Prickly lettuce, Shepherd's purse, Sow cultivate or apply a knockdown herbicide prior to and wheat as directed in thistle/ Milk thistle. Toadrush. Wild applying SPUTNIK, SPUTNIK may also be tankgooseberry, Wild radish, Wild turnip, mixed with certain knock-down herbicides. See the Fallows section of the label Compatibility section of the label. Avoid spray Wireweed/ Hogweed (Polygonum overlap and double spraying on headlands and comers. Refer to the Fallows section of the label (NSW and QLD only) aviculare). Suppression of: Annual ryegrass, Bladder ketmia, for plant-back instructions. Boggabri weed, Caltrop/ vellow vine. Doublegee/ Spiny emex. Dwarf amaranth Phalaris, Pigweed, Wild oats, Wild radish.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS:

Harvest:

NOT REQUIRED WHEN USED AS DIRECTED.

CANOLA, CHICKPEAS, FABA BEANS, FIELD PEAS, LENTILS, LUPINS: DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 6 WEEKS AFTER APPLICATION. SORGHUM: DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 4 WEEKS AFTER APPLICATION

FALLOWS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 6 WEEKS AFTER APPLICATION

www.farmalinx.com



CONTINUED OVERLEAF

sought from Farmalinx Pty Ltd. the breakdown of SPUTNIK in the soil may take longer and further advice should be

L	hese plant-backs are base	isi average seasonal raii	nfall. Under dry conditions,	
]	Barley, Oats, Wheat	4 months	sylnom 3	
]	Mungbeans, Soybeans	shrom &	sdinom 4	
	Following crop after application to fallow	SРUТИІК 0.86 kg/ha	SPUTNIK 1.2 kg/ha	

intended planting date of the following crop thus. Observe the following plant-back intervals between the date of application and the nixed with certain knock-down herbicides. See Compatibility section of the label rain is imminent and prior to germination of weeds. If weeds are present, cultivate or apply a knockdown-herbicide prior to applying SPUTNIK. SPUTNIK may also be tankweeds emerging prior to sowing nominated crops below. Apply to moist soil or when SPUTNIK can be applied to a fallow to provide pre-emergence control of certain

mL/100 L water. stage and sorghum up to 5-7 leaf stage of the crop. Always add Hasten at 500 SPUTNIK can be applied to weeds growing in triazine tolerant canola up to 6-leaf

and treated soil coming on contact with seed. arcc seeders may lead to un-acceptable crop damage due to uneven seeding deput the higher rates, avoid over-lapping sprays and spraying-out corners. Sowing with in particular may cause crop damage from SPUTNIK washed into the seed furrow. At reiatively tiat by the sowing operation. Heavy rain atter application onto ridged soil Pulse crops must be sown to at least 5cm. Apply within 2 days of sowing to soil left Post Sowing Pre Emergence (PSPE)

un-even seeding depth and treated soil coming in contact with seeds. Sowing with disc seeders may lead to un-acceptable crop damage due to shallow,

combination of both press wheels and a covering device such as harrows or chains seed row closure and reduce treated soil falling close to the seed. Do not use a

placement of the seed below the treated soil. The use of press-wheels can help adjacent turrows. Also, poor weed control may occur in deep turrows. Ensure even slow to moderate speed to avoid leaving deep furrows and avoid throwing soil into mm wide and generally placed on 20 cm tine spacing). For crop safety, maintain from SPUTNIK washed into the seed furrow. Crops should preferably be sown with IS minimum-tillage sowing equipment (fitted with knife points or blades less than 12 Heavy rain after application, on ridged soil in particular, may cause crop damage turrows. Crop safety relies on crop seeds germinating below SPUTNIK treated soil. to a week before sowing. The soil should be free of excessive clods, trash and deep Apply to moist soil or when rain is imminent and prior to germination of weeds up

Use only COARSE spray quality according to ASAE S572 definition for standard ground boom in a minimum of 50L/ha. weather conditions when surface temperature inversions can develop. Apply by DO NOT apply when wind speed is less than 3 and greater than 20 km/h or during

canse spray to drift onto nearby susceptible piants/crops, cropping lands or pastures. DO NOT apply under weather conditions, or from spraying equipment, that may NOITACILITATION

beriods, regional use restrictions and safety directions for the tank mix products. Read and follow all label directions, restraints, plant-back periods, withholding

with other products. is advisable that a small scale compatibility test be carried out prior to tank mixing

advice or Farmalinx for current recommendations for control of flax leaf fleabane. It another herbicide such as paraduat ('double knock'). Refer to local agronomist and consideration should be given to cultivation or a subsequent application of mixtures of SPUTNIK plus glyphosate may not control emerged flax leat fleabane application of knockdown herbicides alone, particularly on large weeds. Tank Zome reduction in control may be observed on some weeds compared to the small scale compatibility test be carried out prior to tank mixing with other products. GLYPHO and SIXGUN, PARQUAT, POWERQUAT and PARADAT. It is advisable that a This product is compatible with TRIFLURALINX, SIXGUN, BAZOOKA, WARLORD, **YTIJIBITA9M00**

GENERAL INSTRUCTIONS

SPUTNIK can be used to selectively control weeds in the crops listed in the directions for use table and on fallows prior to establishing crops as directed in Fallows section below. When applied pre-emergence, the product is mainly absorbed via the roots. Weeds buried by cultivation may not be effectively controlled. Heavy rainfall on light soils may cause movement of the herbicide out of the weed seed zone, resulting in reduced weed control. Do not apply to recently burnt stubble/trash. After burning, rainfall or cultivation is required to allow SPUTNIK to reach the soil and not be irreversibly bound by ash. SPUTNIK is primarily incorporated by rainfall. Sufficient rainfall (20 to 30 mm either as rainfall or irrigation) to wet the soil through the weed root zone is necessary within 2 to 3 weeks of application to make the product effective. Weeds may germinate and emerge before being controlled by SPUTNIK however a delay in rainfall for an extended period may result in weeds germinating from depth and surviving. Where soil has a high potential for leaching, heavy rainfal between application and crop emergence may result crop damage from movement of herbicide into the crop seed zone. In flood or furrow irrigation situations, complete and continued activation of the product may not occur due to a thin hand of dry soil on the surface during or after irrigation. Mechanical incorporation after application, using light harrows to incorporate the product into the soil not more than 4 cm deep is required to ensure the irrigation water activates the product. Always apply the product to an even un-ridged seedbed. Dense weed populations and multiple germinations (especially Wild radish) may require both a pre-emergent and post emergent herbicide treatment (only one being SPUTNIK) to give acceptable control.

INTEGRATED WEED MANAGEMENT STRATEGY FOR TT CANOLA

The use of this product in TT canola is subject to the current approved Integrated Weed Management Strategy for the use of triazine herbicides in TT canola". The Strategy encompasses: Integrated Weed Management; the specific management of crop plant volunteers and outcrossing to other plants; and triazine herbicide residue management (agronomic and environment).

Compliance with registered label directions and adoption of the principles outlined in the Strategy will assist with implementation of crop management practices that minimise the development of herbicide resistance in treated weeds; reduce the levels of triazine residues in the environment; and manage volunteer plants and outcrossing. Copies of the Strategy are available from Farmalinx Pty Ltd.
Where specific information and farm management assessment is required to assist

with application of Integrated Weed Management programs on-farm, particularly

To minimise herbicide resistance:

Avoid dry sowing in heavily weed infested paddocks. Wait for weed germination after the opening rains in weedy paddocks. Use a pre-plant knockdown or cultivation. No weeds should be allowed to survive at this stage

in adapting strategies to local conditions, then specific expert advice should

be obtained from an appropriate agronomist, consultant, chemical supplier or

DO NOT use SPUTNIK if the area to be treated had a triazine herbicide applied to

Watch for escapes, especially in paddocks with a long history of Group C herbicide

• DO NOT use Group C herbicides in consecutive years.

Resistant Weeds Reporting

Growers should collect plant or seed samples where weeds that are normally susceptible to atrazine, simazine and terbuthylazine may be resistant, get them tested and seek professional advice.

RESISTANT WEEDS WARNING

FARMALINX SPUTNIK Herbicide is a member of the triazines group of herbicides. FARMALINX SPUTNIK has the inhibitors of photosynthesis at photostem II mode of action. For weed resistance management, FARMALINX SPUTNIK is a Group C herbicide. Some naturally-occurring weed biotypes resistant to FARMALINX SPUTNIK and other Group C herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by FARMALINX SPUTNIK or other Group C herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Farmalinx Pty Ltd accepts no liability for any losses

that may result from the failure of FARMALINX SPUTNIK to control resistant weeds.

DO NOT mix or load within 20 m of any well, dam, intermittent or perennial stream. Partially fill the spray tank with water and maintain agitation while adding the required quantity of SPUTNIK. Agitation should continue after filling and during spraying. If spray is allowed to stand, agitate thoroughly before resuming spraying.