

FARMALINX

Infiltrate

FUNGICIDE

DIRECTIONS FOR USE**RESTRAINTS**

DO NOT apply with equipment carried on the back of the user.

CROP	DISEASE/PEST	STATE	RATE	WHP (DAYS)	CRITICAL COMMENTS
Celery	Septoria leaf spot	Vic, Tas, WA only	200 g/100 L or 2 kg/ha	7	Apply when conditions favour disease and repeat at 7-10 day intervals.
Citrus	Black spot	NSW, Vic, SA, WA only	200 g/100 L plus white oil 250 mL/100 L	7	Apply 6 weeks after copper spray at petal fall and 12 weeks after petal fall. Coastal lowlands and highlands (NSW): apply 8 and 14 weeks after the petal fall copper spray.
	Citrus rust mite	NSW, Vic, SA, WA only	200 g/100 L plus white oil 600 mL/100 L		
	Black spot Citrus rust mite	Qld only	200 g/100 L plus white oil 600 mL/100 L		
Cucurbits e.g.: cucumbers, rockmelons	Downy mildew	All States	200 g/100 L or 2 kg/ha	3	Apply at weekly intervals while conditions favour disease. Spray both sides of leaves thoroughly.
Lettuce	Downy mildew	Vic, Tas, WA only	200 g/100 L or 2 kg/ha	3	Apply when conditions favour disease and repeat at 7-10 day intervals.
Onions	Downy mildew	All States	200 g/100 L or 2 kg/ha	14	Apply at intervals of 7-10 days from early in the life of the crop. If conditions favour disease development consider using a systemic fungicide. Add wetting agent.
Potatoes	Target spot (Early blight) Irish blight (Late blight)	All States	200 g/100 L or 2 kg/ha	1	Apply when conditions favour disease and repeat at 7-10 day intervals.
Tobacco	Blue mould	Vic, WA only	200 g/100 L or 2 kg/ha	7	Apply at weekly intervals commencing 7 days after transplanting and terminating 7 days before first pick. Apply 350 L spray mix per ha after transplanting, increasing to 1350 L on a fully grown crop.
		Qld, NSW, WA only			Apply at weekly intervals terminating 7 days before first pick.
	Frogeye leaf spot Brown spot	Qld, WA only			
Tomatoes	Target spot (Early blight) Irish blight (Late blight) Grey leaf spot (<i>Stemphylium</i>)	All States	200 g/100 L or 2 kg/ha	3	Apply when conditions favour disease and repeat at 7-10 day intervals. In places where diseases are severe commence at seed bed stage.
Watermelon, rockmelon	Anthracnose Gummy stem blight	Qld, WA only	200 g/100 L or 2 kg/ha	3	Apply every 7-10 days depending on weather and disease incidence. First application may be made when vines begin to run.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**WITHHOLDING PERIODS:****Potatoes: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION****Cucurbits, Lettuce, Tomatoes: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION****Celery, Citrus, Tobacco: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION****Onions: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION****GENERAL INSTRUCTIONS****RESISTANT WEEDS WARNING**

Farmalinx INFILTRATE Fungicide is a member of the multi-site group of fungicides. For fungicide resistance management INFILTRATE is a Group M3 fungicide. Some naturally occurring individual fungi resistant to the product and other Group M3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product and other Group M3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Farmalinx Pty Ltd accepts no liability for any losses that result from failure of this product to control resistant fungi.

GROUP M3 FUNGICIDE**Export of treated produce**

Growers should note that MRLs or import tolerances do not exist in all markets for edible produce treated with INFILTRATE. If you are growing edible produce for export, please check with Farmalinx Pty Ltd for the latest information on MRLs and import tolerances before using INFILTRATE.

MIXING

To the required quantity of INFILTRATE add a small amount of water and stir to make a paste. Add further water and mix until a cream results. Pour this into water in spray vat while stirring or with agitators in motion.

RESTRICTIONS

When applying by airblast, use enclosed cab equipment only.

Re-Entry

Do not carry out re-entry activities in citrus orchards for 13 days after treatment, unless wearing gloves and cotton overalls.

COMPATIBILITY

Ensure that water is of suitable quality for mixing with agricultural pesticides. Poor quality water can cause physical incompatibility between some products which may result in crop damage in some circumstances. INFILTRATE is compatible with most commonly used insecticides and fungicides including Bayfidan® 250 EC and Dipterex® 500 SL. INFILTRATE is not compatible with lime sulphur and other alkaline materials. When mixing sprays, except when adding oil, add insecticides last. Do not mix concentrates together but add each to the spray tank separately. The compatibilities given are based on world wide experience and are considered correct. However, as changes in climatic conditions can alter the sensitivity of plants to mixtures of sprays, we accept no responsibility for the behaviour of such mixtures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or watercourses with this product or used containers.

STORAGE AND DISPOSAL**Containers**

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

NOTICE:

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Farmalinx Pty Ltd or under abnormal conditions. Farmalinx Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

© Registered Trade Marks

Bags

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight.

Single-rinse or shake remainder into spray tank. Do not dispose of undiluted chemicals on site. Puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Harmful if inhaled. Avoid inhaling dust. Avoid contact with skin. When opening the container and preparing spray, wear cotton overalls (over normal clothing) buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves and a half face-piece respirator with dust cartridge or canister. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. Avoid giving alcohol.

SDS

Additional information is listed in the safety data sheet (SDS). A safety data sheet for Farmalinx INFILTRATE Fungicide is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 9389 2455.

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FARMALINX
Infiltrate

FUNGICIDE

ACTIVE CONSTITUENT: 700 g/kg PROPINEB

GROUP M3 FUNGICIDE

For the control of blue mould of tobacco, target spot (early blight) and late blight (Irish blight) of potatoes and tomatoes, grey leaf spot of tomatoes, downy mildew of cucurbits and onions, black spot of citrus and certain other diseases as specified in the Directions for Use table.

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.

FARMALINX Pty Ltd ABN 95 134 353 245
Level 25, Suite 2506, Tower 2, 101 Grafton Street,
Bondi Junction NSW 2022

farmalinx

farmalinx
Partners in Agriculture

FARMALINX Pty Ltd ABN 95 134 353 245

Level 25, Suite 2506, Tower 2,

101 Grafton Street, Bondi Junction NSW 2022

T: 02 9389 2455 F: 02 9389 2844

Website: www.farmalinx.com

APVMA Approval NO.: 83339/108083