


Creepers 800 WG

farmalinx



DANGEROUS POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FARMALINX

Creepers 800 WG

FUNGICIDE

ACTIVE CONSTITUENT: 800 g/kg CARBENDAZIM

GROUP 1 FUNGICIDE

A broad spectrum, systemic fungicide for the control of fungal diseases in various crops as per the Directions for Use table.

farmalinx *Partners in Agriculture*

FARMALINX Pty Ltd ABN 95 134 353 245
Level 25, Suite 2506, Tower 2, 101 Grafton Street, Bondi Junction NSW 2022
T: 02 9389 2455 F: 02 9389 2844 W: www.farmalinx.com

NET CONTENTS
15 kg



Creepers 800 WG

farmalinx

DIRECTIONS FOR USE

1. Tree Crops

RATE: This table shows rates for dilute spraying. For concentrate spraying, refer to the Mixing/Application section.			CRITICAL COMMENTS For all uses in this table: Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. For concentrate spraying, DO NOT use at rates greater than 5 times the dilute spraying rate		
CROP	DISEASE	STATE	RATE	WHP (DAYS)	CRITICAL COMMENTS
Macadamia nuts	Macadamia Husk Spot (<i>Pseudocercospora</i> spp.)	All States	31 g/100 L plus a wetting agent (at 100 mL/100 L)	14 (H) 28 (G)	Apply at 5 and 8 weeks after main flowering - Stage 2 anthesis (white flowering stage). Remove any fallen nuts from under trees prior to spraying. DO NOT apply more than 2 applications of CREEPER 800 WG per season.

2. Other Crops

CROP	DISEASE	STATE	RATE	WHP (DAYS)	CRITICAL COMMENTS
Chickpeas, faba beans, lentils and vetch	Chocolate Spot (<i>Botrytis fabae</i>), Grey Mould (<i>Botrytis cinerea</i>)	All States	313 g/ha	4 weeks (H, G)	Apply a maximum of two consecutive applications at 14 day intervals. Apply a minimum of 100L of water per hectare.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIOD:
MACADAMIA NUTS: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.
DO NOT ALLOW LIVESTOCK TO GRAZE VEGETATION IN TREATED PLANTATIONS FOR 4 WEEKS AFTER APPLICATION.
DO NOT CUT VEGETATION IN TREATED PLANTATIONS FOR STOCK FEED FOR 4 WEEKS AFTER APPLICATION.
CHICKPEAS, FABA BEANS, LENTILS AND VETCH: DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION.
DO NOT GRAZE OR CUT FOR STOCKFEED FOR 4 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

MIXING/APPLICATION
SHAKE WELL BEFORE USE. FARMALINX CREEPER 800 WG FUNGICIDE is a water dispersible granule to be mixed with water for application as a spray. Add the required quantity of CREEPER 800 WG to a partly filled spray tank and agitate. Complete filling while agitating. Ensure the spray mixture is properly agitated before restarting after stoppage.

ADDITION OF WETTING AGENT
The addition of a wetting agent eg. Activator® Surfactant to spray mixture improves the distribution of the spray on hard to wet plants.

APPLICATION
Dilute Spraying (Tree crops only)
Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive runoff. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. Add the amount of product specified in the Directions for Use table for each 100 L of water. Spray to the point of run-off. The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate Spraying (Tree crops only)
Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume. Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate. The mixing rate for concentrate spraying can be calculated in the following way:
EXAMPLE ONLY
1. Dilute spray volume as determined above: For example 1500 L/ha
2. Your chosen concentrate spray volume: For example 500 L/ha
3. The concentration factor in this example is: 3X (i.e. 1500 L ÷ 500 L = 3)
4. If the dilute label rate is 10 mL/100 L, then the concentrate rate becomes 3 x 10, that is 30 mL/100 L of concentrate spray.
The chosen spray volume, amount of product per 100 L of water, the sprayer set up and operation may need to be changed as the crop grows. For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

FUNGICIDE RESISTANCE WARNING

FARMALINX CREEPER 800 WG FUNGICIDE is a member of the benzimidazole group of fungicides. For fungicide resistance management CREEPER 800 WG is a Group 1 fungicide. Some naturally occurring fungal populations resistant to CREEPER 800 WG and other Group 1 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. The resistant fungi will not be controlled by CREEPER 800 WG and other Group 1 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Farmalinx Pty Ltd accepts no liability for any losses that may result from the failure of CREEPER 800 WG to control resistant fungi.

EXPORT OF TREATED PRODUCE

Growers should note that suitable MRLs or import tolerances may not be established in all markets for produce treated with CREEPER 800 WG. If you are growing produce for export, please check with Farmalinx for the latest information on MRLs and import tolerances BEFORE using CREEPER 800 WG.

COMPATIBILITY

CREEPER 800 WG is compatible with many insecticides and fungicides, however to be sure of compatibility with any particular product check with your local dealer first.

PRECAUTIONS

Keep away from the foodstuffs.
RE-ENTRY PERIOD
DO NOT allow entry into treated areas until the spray has dried unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET CROPS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.
PROTECTION OF LIVESTOCK
CREEPER 800 WG is of low toxicity to bees.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life with long lasting effects. DO NOT contaminate wetlands or watercourses with this product or used containers.
STORAGE AND DISPOSAL
Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Shake and empty contents into spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulation. Do not burn empty containers or product.

SAFETY DIRECTIONS

Very dangerous, particularly the concentrate. Poisonous if absorbed by skin contact, inhaled or swallowed. May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. Do not inhale vapour or spray mist. When opening the container, and preparing the spray wear elbow-length chemical resistant gloves. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 13 11 26.
WARNING: Contains carbendazim which causes birth defects and irreversible male infertility in laboratory animals. Avoid contact with carbendazim.

SDS

Additional information is listed in the safety data sheet (SDS). A material safety data sheet for Farmalinx Creeper 800 WG Fungicide is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 9389 2455.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Farmalinx Pty Ltd or under abnormal conditions. Farmalinx Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

May damage fertility. May damage the unborn child.
Obtain special instructions before use.
Do not handle until all safety precautions have been read and understood.
If exposed or concerned: Get medical advice/attention.

IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE

SPECIALIST ADVICE IN AN EMERGENCY ONLY
0417 653 654
ALL HOURS – AUSTRALIA WIDE



Batch No.
Date of Manufacture