CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FARMALINX



INSECTICIDE

ACTIVE CONSTITUENT: 25 g/kg CLOTHIANIDIN

GROUP INSECTICIDE

A combined slow release fertiliser and insecticide treatment, for the control of leaf eating insects in Eucalypt seedlings

GUARANTEED FERTILISER ANALYSIS 16% TOTAL NITROGEN 12.5% NH₄ nitrogen 3.5% NO₃ nitrogen 25% PHOSPHATE P₂O₅

21% P₂O₅ water soluble 6% POTASSIUM oxide K₂O water soluble 1% MAGNESIUM Mg0 0.7% water soluble Mg0 2.5% SULPHUR (S) 2.0 % water souble S

farmalinx Partners in Agriculture

FARMALINX Pty Ltd ABN 95 134 353 245 Level 25, Suite 2506, Tower 2, 101 Grafton Street, Bondi Junction NSW 2022 T: 02 9389 2455 F: 02 9389 2844 W: www.farmalinx.com

CROP	PEST	RATE	CRITICAL COMMENTS
Eucalypt	Leaf feeding	20 g/seedling	Apply at planting using an enclosed planting applicator into a
seedlings	beetles:		planting tube or at or after planting by subsoil placement.
from planting	Including:		
to 1 year old	Paropsisterna		Planting tube: Place 20 g of Sucker G at the bottom of the planting
	agricola,		tube using an enclosed planting applicator, then add the seedling,
	Cadmus		insert the planting tube in the ground and open to release the
	australis,		contents.
	Liparetrus		
	jenkinsi,		Subsoil placement: Using a long handled spade make a cut 2 cm
	Heteronyx		from the root ball 10 cm deep. Apply 20 g of product using an
	<i>dimidiata</i> and		enclosed planting applicator then close the cut.
	other		
	Heteronyx		DO NOT apply SUCKER G directly to the root ball area as this may
	species.		affect plant growth. The speed of control from this application
			depends on how fast the product is taken up by the seedling, which
			needs to be actively growing. This will be assisted by post-planting
			rainfall or watering of at least 10 mm. It is recommended that
			application is before or during the main annual rainfall period. If
			severe attack takes place prior to the chemical becoming effective
			in the plant, a knockdown insecticide may be required to give short
			term protection.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

Farmalinx Sucker G Insecticide is a specially coated fertiliser which provides controlled release of essential nutrients to the seedlings and provides protection against attack from the species listed in the Directions for Use table. Clothianidin is a systemic insecticide which is absorbed into the plants via the roots and is translocated through the seedling as it grows.

biotypes resistant to SUCKER G and other Group 4A insecticides may exist through normal genetic variability

in any insect population. The resistant individuals can eventually dominate the insect population if SUCKER

G or other Group 4A insecticides are used repeatedly. The effectiveness of SUCKER G on resistant individuals

could be significantly reduced. Since the occurrence of resistant individuals is difficult to detect prior to

use, Farmalinx Pty Ltd accepts no liability for any losses that may result from the failure of this product to

control resistant insects. SUCKER G may be subject to specific resistance management strategies. For further

INSECTICIDE RESISTANCE WARNING

For insecticide resistance management Farmalinx SUCKER G GROUP 4A INSECTICIDE Insecticide is a Group 4A insecticide. Some naturally occurring insect



APVMA Approval No. 86898/116932

information contact your local supplier, Farmalinx Pty Ltd representative or local agricultural department agronomist. IN A TRANSPORT EMERGENCY **DIAL 000**

SAFETY DIRECTIONS

FIRST AID If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia Tel. 131126; New Zealand 0800 764 766.

SDS Additional information is listed in the safety data sheet (SDS). A safety data sheet for Farmalinx SUCKER G Insecticide is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 9389 2455.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Farmalinx Pty Ltd or under abnormal conditions. Farmalinx Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.







ROTECTION OF LIVESTOCK

angerous to bees and will kill bees foraging in the crop to be treated. sks to non-target insects - Clothianidin may have adverse effects on some non-target beneficials, in rticular where IPM is practiced, to foliage dwelling predators.

ROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

nis product is highly toxic to aquatic invertebrates. **DO NOT** contaminate streams, rivers or waterways with chemical or used containers.

FORAGE AND DISPOSAL

core in the closed, original container in a dry, cool, well-ventilated area, away from direct sunlight. Triple or eferably pressure rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted nemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. not recycling, break, crush or puncture and deliver empty packaging to an approved waste management cility. If an approved waste management facility is not available bury the empty packaging 500 mm below surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable getation and tree roots, in compliance with relevant Local, State or Territory government regulations. **NOT** burn empty containers or product.

refillable containers: Empty contents fully into application equipment. Close all valves and return to point supply for refill or storage.

Poisonous if swallowed. Wash hands after use.



Batch No.

Date of Manufacture

