

DIRECTIONS FOR USE

Restraints

DO NOT spray when rain seems likely to occur with 4 hours.

DO NOT spray when weeds are wet with dew or rain or under stress from drought, low soil fertility, extreme cold or water logging

DO NOT spray outside recommended crop growth as crop damage may result.

DO NOT apply to crops or pastures with clover, lucerne or medics.

DO NOT apply to legume pastures.

DO NOT apply to plants suffering from stress.

Crop & Situation	Weeds	State	Rate/ha	Critical Comments
Barley, Commercial Rye, Oats, Triticale, Wheat From early tillering stage (Zadoks 21)	Capeweed, Charlock, Common Ice- Plant, seedling Docks, Doublegee (Three cornered Jack, Spiny Emex) Hexham Scent (Melilotus), Mustards, Solider Thistle, Sorrel Seedlings, Tares, Tree Hogweed, Turnip Weed (<i>Rapistrum rugosum</i>), Vetches, Ward's Weed, Wild Radish, Wild Turnip (<i>Brassica</i> tournefortii), Wireweed (Hogweed)	All States	1L	Spray when most weeds are in the 2 to 4 leaf stage. In some instances suppression only of Wireweed (Hogweed), Capeweed and Wild Radish will occur.
	Seedling Cotton Thistle, Fat-Hen	Tas		
Barley, Commercial Rye, Oats, Triticale, Wheat From early tillering stage (Zadoks 21) until the fully tillered stage and before jointing occurs (Zadoks 30)	Cotton Thistle to bud formation	Tas	1.7 L Spray when most weeds are still in the rosette or seedling stage. N.B. Wild radish may need to be resprayed (with 2.4-D or	
	Mintweed, New Zealand Spinach	Qld, NSW		
	Capeweed, Charlock, Chickweed, Climbing Buckwheat (Black Bindweed), Clover, Common Ice-Plant, Seedling Docks, Doublegee (Three cornered Jack, Spiny Emex), Fat-Hen, Hexham Scent (Melilotus), Seedling Hoary Cress, Mouse-eared Chickweed, Mustards, Narrow-leaved Plantain, Saffron Thistle, Spurry, Swinecress, Tares, Tree Hogweed, Turnip Weed (<i>Rapistrum rugosum</i>), Variegated Thistle, Vetch, Wild Radish, Wild Turnip (<i>Brassica tournefortil</i>), Wireweed (Hogweed)	Qld, NSW, Tas, WA, Vic		
		SA	1.4 L	
Barley, Commercial Rye, Triticale From early tillering stage (Zadoks 21) until the fully tillered stage before jointing occurs (Zadoks 30)	Volunteer Peas	SA	1 L plus 60 mL/ha Spalding Dicamba 500 Selective Herbicide	Apply when most peas are 5-7cm high.
Oats, Wheat From early tillering stage (Zadoks 21) until the fully tillered stage before jointing occurs (Zadoks 30)			1 L/ha plus 120 mL/ha Spalding Dicamba 500 Selective Herbicide	

storage, handling or use. accepts no liability for any loss or damage arising from incorrect Farmalinx Pty Ltd or under abnormal conditions. Farmalinx Pty Ltd to label instructions, or under off-label permits not endorsed by burpose, express or implied, extends to the use of product contrary statutory warranties) of merchantability or fitness for a particular conditions of use. No warranty (other than non-excludable label when used in accordance with directions under normal description and is reasonably fit for the purposes stated on the Seller warrants that the product conforms to its chemical

CONDITIONS OF SALE:

SDS

CAUTION

02 9389 2455 from Farmalinx Pty Ltd on request. Call Customer Service on A safety data sheet for Farmalinx SAVIOR Herbicide is available Additional information is listed in the safety data sheet (SDS).


Centre (Telephone Australia 13 11 26).

If poisoning occurs, contact a doctor, or Poisons Information **GIA TZRIT**

use, wash gloves, face shield and contaminated clothing. arms and face thoroughly with soap and water. After each day's After use and before eating, drinking or smoking, wash hands, If product on skin, immediately wash area with soap and water. PVC gloves. If product in eyes, wash out immediately with water. buttoned to the neck and wrist, washable hat and elbow length or goggles. When using the prepared spray wear cotton overalls and wrist, washable hat, elbow-length PVC gloves, face shield sud preparing spray wear cotton overalls buttoned to the neck exposure may cause allergic disorders. When opening container Will irritate the eyes and skin. DO NOT inhale vapour. Repeated Harmful if absorbed by skin contact, inhaled or swallowed. SAFETY DIRECTIONS

353 Street 134 afton Ltd Tower FARMALINX F Level 25, Suite 25 Bondi Junction MS Certain Bi

KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING ACTIVE CONSTITUENTS: 340 g/L MCPA present as the 80 g/L DICAMBA present as the dimethylamine salt HERBICIDE For the Control of C Cereals, Pastures, 1 Directions for Use. GROUP



NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED **UNDER APPROPRIATE LEGISLATION**

WITHHOLDING PERIODS

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION HARVEST WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED

www.farmalinx.com



CONTINUED OVERLEAF

chemical or used containers.

DO NOT contaminate streams, rivers or waterways with the DO NOT use container for any other purpose.

TNAMNORIVNA DNA

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS

such as cotton, tobacco, vines, lupins, fruit trees and ornamentals. Avoid spray drift and vapour movement onto susceptible crops crops, cropping lands or pastures.

equipment, that may cause drift onto nearby susceptible plants/ DO NOT apply under weather conditions, or from spraying use and must not be used for spraying horticultural crops.

in use. All spray equipment must be thoroughly washed out after contact with their roots. Keep container sealed (airtight) when not in locations where the chemical may be washed or moved into DO NOT apply this product on or near desirable trees or plants or NOT give a fine droplet size. DO NOT use in high wind.

Minimise spray drift by using low pressure and nozzles which DO

Spray equipment must be calibrated accurately prior to working

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET

GENERAL INSTRUCTIONS

Add the required amount of SAVIOR directly to the tank and mix well

Local, State or Territory government regulations. Do not burn

desirable vegetation and tree roots, in accordance with relevant

specifically marked and set up for this purpose clear of waterways,

embty packaging 500mm below the surface in disposal pit approved waste management facility is not available bury the

backaging to an approved waste management facility. If an

If not recycling, break, crush or puncture and deliver empty

kepiace cap and return clean container to recycler or designated

tank. DO NOT dispose of undiluted chemicals on-site. If recycling,

Triple rinse containers before disposal. Add rinsings to spray

fertilizers. DO NOT store for prolonged periods in direct sunlight.

area away from children, animals, food, feedstuffs, seed and

Store in the closed, original container in a dry, cool, well-ventilated

empty containers or product.

collection point.

STORAGE AND DISPOSAL

APPLICATION

For Boom Application: Apply by a properly calibrated boom spray using not less than 50 litres of water per hectare unless

indicated otherwise in Critical Comments column. For Aerial Application: Apply at least 25 litres total spray volume per hectare. The aircraft should fly as low as practical under the prevailing conditions to minimise drift.

For High Volume Application: Apply minimum 1500 litres per hectare.

For Knapsack Application: A 15 litre knapsack treats 150 square metres

NOTE: Spray when weeds are young and actively growing. A temporary wilting may be evident in crops after application. If seasonal conditions are unusually late or dry seek further advice as damage may result from spraying under these conditions. Growers should seek advice before spraying recently released cereal varieties. There is nil tolerance of pea seed contaminates in cereal grain and less than 100% control of Volunteer Peas could result in dockage

COMPATABILITY

SAVIOR is compatible with flowable terbutryn.

RESISTANT WEEDS WARNING

Farmalinx SAVIOR Herbicide is a GROUP HERBICIDE member of the Benzoic acid and

Phenoxy groups of herbicides. SAVIOR has the disruptors of plant cell growth mode of action. For weed resistance management SAVIOR is a Group I Herbicide. Some naturally occurring weed biotypes resistant to SAVIOR and other Disruptors of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by SAVIOR or other Group I Herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Farmalinx Pty Ltd accepts no liability for any losses that may result from the failure of SAVIOR to control resistant weeds.



FARMALINX Pty Ltd ABN 95 134 353 245 Level 25, Suite 2506, Tower 2, 101 Grafton Street, Bondi Junction NSW 2022 T: 02 9389 2455 F: 02 9389 2844 Website: www.farmalinx.com

APVMA Approval No. 88327/120869