Grasses

Bahia Grass (Paspalum notatum) Suppression and Seedhead Control

AUTHORISED UNDER APPROPRIATE LEGISLATION.

Grasses

Bahia grass (Paspalum

notatum) suppression

and seedhead control

Winter Grass

Poa annua

Rate

/100m²

Rate

non-ionic surfactant (such as Agral)

4 to 10 mL

/ha

2 to 4 L plus 75

Herbicide plus 0.25 to 0.42% v/v

(such as Agral

Spray Adjuvant)

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS

DO NOT GRAZE TREATED TURF/LAWN, OR FEED TURF/LAWN CLIPPINGS FROM ANY TREATED AREA TO POULTRY OR LIVESTOCK.

Liquid Turf

mL/ha Monument

/ha

0.4 L to 1 L

Winter Grass (Poa annua) Management

Situation

Golf course fairways,

Situation

Common couch. Durban

grass, Hybrid couch.

Qld Blue couch, Zoysia

only in aolf course long

roughs, lawns, median

WITHHOLDING PERIODS

strips, ovals, parks,

roadsides, general

grassed areas.

Golf course greens

Critical Comments

Critical Comments

Apply during summer when Bahia grass

DO NOT apply more than 2 consecutive

seedheads in established couch areas.

Refer to Application section for detailed

applications of Monument Liquid per season

See instructions under Bahia grass (Paspalum notatum) Management for control of Bahia Grass

Important: Refer to the Monument Liquid label for specific directions on mixing, avoiding spray drift, sprayer cleanup and protection of nontarget crops and the environment

See instructions under Winter Grass (Poa annua) Management for

produces seedheads

information.

reduction of Poa annua swards in Bent Grass golf greens. See Golf course greens section above for application

recommendations

/100m²

20 to 40 mL plus

0.75 mL/100 m² of

0.25 to 0.42% v/v

non-ionic surfactant

Monument Liquid plus

Bingo ME 120

DIRECTIONS FOR USE

Restraints

DO NOT apply to turf recovering from wear, renovation or turf affected by disease, waterlogging or other stresses. DO NOT use where growth regulatory effects are likely following the application of a fungicide or any other material. DO NOT apply to turf going into or coming out of dormancy. DO NOT apply if rainfall is expected within 1 hour, and DO NOT irrigate for at least 1 hour after application. DO NOT apply to Kentucky Bluegrass as damage may occur.

Turf Growth Management

		Rate		
Situation	Grasses	/ha	/100m ²	Critical Comments
Golf course and bowling greens	Bent grass Bent Grass/ Winter Grass <i>(Poa annua</i>) mixed sward	0.2 to 1 L	2 to 10 mL	 Apply BING0 ME 120 at least 6 hours before or after mowing and when turf is actively growing. Use the lower rates on turf with moderate growth rates where a lower level of growth reduction is required or when used in a program. Apply rates towards the highest end of the rate range when growth is vigorous or when longer periods of 4 to 6 weeks regulation are required. Application made to turf growing under cool conditions or to turf with low nutrient levels or where growth rates are low may result in discolouration. Well-maintained lawns and turf surfaces are those that receive at least weekly mowing and good levels of water and fertility are maintained. Refer to Application Section for detailed information.
	Hybrid couch	0.15 to 0.6 L	1.5 to 6 mL	
Golf course fairways	Carpet grass Kikuyu	1.6 to 4 L	16 to 40 mL	
	Common couch	0.8 to 1.6 L	8 to 16 mL	
	Couch varieties: Greenlees Park, Santa Ana, Wintergreen	0.6 to 1.2 L	6 to 12 mL	
	Fescue, Perennial Ryegrass	1.6 to 3 L	16 to 30 mL	
Golf course short roughs, well-maintained lawns and turf surface	Bahia Grass, Bent grass, Fescue, Perennial Ryegrass	2 to 4 L	20 to 40 mL	
	Buffalo grass Common couch, Hybrid couch	1 to 2 L	10 to 20 mL	
	Kikuyu	1.6 to 6 L	16 to 60 mL	
Golf course long roughs, parks, ovals, lawns, general grassed areas	Bent Grass, Fescue, Perennial Ryegrass	2 to 6 L	20 to 60 mL	
	Buffalo Grass Common Couch, Hybrid couch	1 to 4 L	10 to 40 mL	
	Kikuyu	2 to 8 L	20 to 80 mL	
Bunker facing, path and garden edging, footpath and kerbside edging, fencelines, around trees.	Bent Grass, Common couch, Perennial Ryegrass	4 to 6 L	40 to 60 mL	Apply BINGO ME 120 at least 6 hours before or after mowing and when turf is actively growing.
	Buffalo grass	4 L	40 mL	On difficult to mow areas apply BINGO ME 120 to actively growing turf and mow 3 to 4 days after application. Apply in a strip about 25 to 50 cm wide.
	Couch varieties: Greenlees Park, Wintergreen, Santa Ana			
	Kikuyu	6 to 8 L	60 to 80 mL	Use the lowest rate of BINGO ME 120 around bunker edges and the higher rates in the other situations.
				Application made to turf growing around cool conditions or to turf with low nutrient levels or where growth rates are low may result in discolouration.

www.farmalinx.com

CONTINUED OVERLEAF

.bnud.

and 3 bar pressure. Boom height should not exceed 50 cm above the coarse [Turbo Teejet or AIXR Teejet (11004 or 11005)] nozzles, at 8 km/h application volume should be 200 to 500 L/ha. For best results use

placement as uniformly as possible onto leaves and into crowns. Ideal Bahia Grass Suppression and Seedhead control: Ensure product .bnud.

hr and 3 bar pressure. Boom height should not exceed 50 cm above the coarse [Turbo Teejet* or AIXR Teejet* (11004 or 11005)' nozzles, at 8 km/ 500 L/ha. For best results use medium [XR Teejet* (11004 to 11008)] to onto leaves and into crowns. Ideal application volume should be 200 to All Other Turf Areas: Ensure product placement as uniformly as possible not exceed 50 cm above the ground.

or 11005)] nozzles, at 5 km/hr and 3 bar pressure. Boom height should Teejet* (11004 or 11005)] to coarse [Turbo Teejet* or AIXR Teejet* (11004 Volume should be 350 to 500 L/ha. For best results use medium [XR uniformly as possibly onto leaves and into crowns. Ideal application Golf Course and Bowling Greens: Ensure product placement as

 m^2 of turf. Preferably use flat fan nozzles and ensure even coverage is 120 to a known area of turt. Apply between 2 and 10 L of water / 100 Knapsack: It is important to apply the correct quantity of BINGO ME semulov notresilqqa.

'sspifi

out spraying equipment, mixing and measuring utensils with water onto streams or storm water drains with concentrate or used container. Wash Dangerous to fish larvae and oysters. DO NOT contaminate dams, rivers, PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

until spray has dried.

until spray has dried. DO NOT allow the public to enter treated areas Re-entry Period: D0 NOT enter treated areas, without protective clothing **PRECAUTIONS**

the broadleaf weed herbicides.

allowing at least 10 days between an application of BINGO ME 120 and to apply BINGO ME 120 and the broadleaf weed herbicides separately, turf with herbicides other than those indicated above, it is recommended after application of a tank mix. For the treatment of broadleaf weeds in However, some transient discolouration of the treated turf may occur dicamba or MCPP/dicamba.

Bromoxynil, MCPP ester/bromoxynil/dicamba, MCPP/2,4-D amine/ 2,4-D amine, bromoxynil, MCPA/bromoxynil/dicamba, MCPA ester/ containing the following chemicals:

BINGO ME 120 is compatible with some formulations of products For common Couch, Hybrid Couch, and Kikuyu turf grasses only:

120 around other materials, test on a small scale basis first. concrete, granite, marble or other types of stone. Before using BINGO ME

from FARMALINX Pty Ltd on request. Call Customer Service on 02 9389 asta sheet for FAMALINX BINGO ME 120 Growth Regulator is available Ytəts A. (SDS) teacher data sheet vata sheet (SDS). A safety TEERS ATAD YTEERS

> Telephone Australia 13 11 26, New Zealand 0800 764 766. If poisoning occurs, contact a doctor or Poisons Information Centre. **DIA TZAI**

gloves, face shield or goggles and contaminated clothing. face shield or goggles. Wash hands after use. After each day's use, wash neck and wrist, elbow-length nitrile or PVC gloves, a washable hat and spray and using prepared spray, wear cotton overalls buttoned to the it out immediately with water. When opening the container, preparing Will irritate the eyes. Avoid contact with eyes. If product in eyes, wash SNOITCIAN DIRECTIONS

Jonduct.

ABN (101 (

Ltd Tower 2022

FARMALINX Pty L1 Level 25, Suite 2506, Tc Bondi Junction NSW 20

rmalinx

Territory government regulations. DO NOT bury empty containers or vegetation and tree roots, in compliance with relevant local, State or marked and set up for this purpose clear of waterways, desirable packaging 500 mm below the surface in a disposal pit specifically an approved waste management facility is not available, bury the empty

storage, handling or use. Pty Ltd accepts no liability for any loss or damage arising from incorrect

μραςτική πεσατινείν οπ spray penetration and even coverage at iow higher cut turf (>15 mm) a significant spray shielding effect can occur, and even coverage of the turf leaves and penetration to the crowns. In fan nozzles. Application volume should be adequate to ensure thorough Application Volume: Apply in 100 to 500 L/ha water using suitable flat overlapping sprays.

Ensure even application in order to achieve even growth reduction. Avoid Application

Bahia Grass (Paspalum notatum) Management

Different grasses vary in their sensitivity to BINGO ME 120 and Monument Liquid. Bahia Grass is very senstivie to the combination resulting in control of seedheads as well as vegetative growth suppression. Apply the combination as soon as seedheads start to form or when it becomes a regular mowing intervention issue, normally during October to January. Apply at 4 week intervals after mowing

Couch Overseeding

An application of BINGO ME 120 to couch will enhance the establishment of cool season turf grasses. BINGO ME 120 is foliarly absorbed, so germination and seedling growth is not affected. Apply BINGO ME 120 before verticutting, scalping, spiking or other similar operations to the couch. Apply BINGO ME 120 1 to 5 days before seeding.

Mixing

BINGO ME 120 is a micro-emulsion concentrate (ME) with minimal odour, as the product does not contain petroleum solvents. BINGO ME 120 mixes completely with water and may be tank mixed with many commonly used fungicides and liquid fertilisers. Add the required quantity of BINGO ME 120 directly to a spray tank containing 2/3 (two thirds) of the required volume. Add the rest of the water and ensure the mix is thoroughly agitated before application.

BINGO ME 120 can be tank mixed with Heritage MAXX Turf Fungicide, Banner MAXX Turf Fungicide, Daconil Weather Stik Turf Fungicide and Monument Liquid.

at. 042% v/v or 420 mL/100 L spray mix. Use surfactants (1000 g ai/L control when tank mixing with Monument Liquid, use Agral (600 g ai/L) Non-ionic surfactants: For Bahia Grass suppression and seedhead

Farmalinx BINGO ME 120 Growth Regulator should be applied to actively

growing turf, at least 6 hours before or after mowing. Good mowing

management leading up to application is required to achieve a good

quality turf surface and to obtain the best results from BINGO ME 120.

The recommended use rates are designed to give approximately a 20

to 50% reduction in clippings over a 2 to 6 week period. Full growth

regulation by BINGO ME 120 begins about 3 to 5 days after application.

In practise, the amount of growth reduction achieved will vary between

situations due to environmental conditions and management practices.

Therefore, to some extent application rates will need to be adjusted to

match growing conditions, management practices and the amount of

Repeat applications can be made as soon as turf resumes growth or

more suppression is required. Areas treated with BINGO ME 120 should

continue to receive regular maintenance practises, including irrigation, fertilisation, and weed, disease and insect control where necessary, as

Different grasses vary in their sensitivity to BINGO ME 120. Since Poa

annua is more sensitive to BINGO ME 120 than Bent grass, BINGO ME

120 can be used to reduce the competitiveness of Poa annua in Bent

grass golf greens. BINGO ME 120 can be integrated into existing Poa

growth suppression of Poa annua, but not Bent Grass) applied during

peak growth periods may enable slow transition in the golf green sward

composition away from Poa annua and towards a higher percentage of

annua control programs. Low rates of BINGO ME 120 (that provide

growth regulation required by the turf manager.

Winter Grass (Poa annua) Management

recommended for quality turf.

Bent grass.

BINGO ME 120 at normal dilution rates, will not stain brass, bronze, Application Near and Around Monuments and Hardscape Materials.

formulations) at 0.25% v/v or 250 mL/100 L spray mix.

JA2092ID & JDA90T2

GENERAL INSTRUCTIONS

Growth Regulation

for appropriate disposal to an approved waste management facility. If If not recycling, break, crush or puncture and delivery empty packaging return clean containers to recycler or designated collection point. NOT dispose of undiluted chemicals on site. If recycling, replace cap and Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT store for prolonged periods in direct sunlight.

Store in the closed, original container in a cool, well-ventilated area. DO

endorsed by Farmalinx Pty Ltd or under abnormal conditions. Farmalinx of product contrary to label instructions, or under off-label permits not fitness for a particular purpose, express or implied, extends to the use (other than non-excludable statutory warranties) of merchantability or accordance with directions under normal conditions of use. No warranty and is reasonably fit for the purposes stated on the label when used in Seller warrants that the product conforms to its chemical description

CONDITIONS OF SALE:

aid **CAUTION** Keep out of reach of children Read safety directions before opening or using IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER Read thoroughly before opening or using this product. Stem growth of Grass management and as a ACTIVE CONSTITUENT: 120 g/L TRINEXAPAC-ETHYL TOR For the reduction of Leaf and species and as an aid to Turf i in Winter grass management. **GROWTH REGULA —**] **A**

