



## DIRECTIONS FOR USE RESTRAINTS

DO NOT apply in Tasmania in autumn or summer.  
DO NOT apply in southeast Queensland unless in July or August.  
DO NOT apply in Mary Burnett in April, May, October or November.  
DO NOT apply by vertical sprayer.  
DO NOT apply by fixed wing aircraft.  
DO NOT apply if heavy rains or storms are forecast within 3 days.  
DO NOT irrigate to the point of runoff for at least 3 days after application.  
DO NOT apply to waterlogged soil.  
DO NOT apply more than 205 kg Farmalinx Weapon Herbicide (182 kg terbuthylazine) per day by air, or 78 kg Farmalinx Weapon Herbicide (68.25 kg terbuthylazine) per day by ground boom.

### Spray Drift Restraints:

Specific definitions for terms used in this section of the label can be found at [apvma.gov.au/spraydrift](http://apvma.gov.au/spraydrift).  
DO NOT allow bystanders to come into contact with the spray cloud.  
DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone table/s below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.  
DO NOT apply unless the wind speed is between three and 20 kilometres per hour at the application site during the time of application.  
DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DO NOT apply by a boom sprayer unless the following requirements are met:

- Spray droplets are not smaller than a COARSE spray droplet size category.
- Minimum distances between the application site and downwind sensitive areas are observed (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for boom sprayers').

### Buffer zones for boom sprayers

Application rate	Boom height above the target canopy	Mandatory downwind buffer zones		
		Bystander areas	Natural aquatic areas	Vegetation areas
Up to 5.2 kg/ha	0.5 m or lower	50 metres	160 metres	70 metres
3.5 kg/ha	0.5 m or lower	50 metres	100 metres	50 metres
	1.0 m or lower	50 metres	375 metres	140 metres

DO NOT apply by aircraft unless the following requirements are met:

- For application by helicopter only, not for application by fixed-wing aircraft.
- Spray droplets are not smaller than a COARSE spray droplet size category.
- For maximum release height above the target canopy of 3 metres or 25 per cent of wingspan or 25 per cent of rotor diameter, whichever is the greatest, minimum distances between the application site and downwind sensitive areas are observed (see the following table titled 'Buffer zones for aircraft')

### Buffer zones for aircraft

Type of aircraft	Mandatory downwind buffer zones		
	Bystander Areas	Natural aquatic areas	Vegetation areas
Helicopter	130 metres	800 metres	400 metres

CROP	SITUATION	WEEDS	RATE	CRITICAL COMMENTS
Pinus radiata	Pre-planting	African daisy Common Storksbill Chickweed Fleabane Kikuyu grass <sup>1</sup> Phalaris Red sorrel <sup>2</sup> Silver grass Soft brome Spear thistle Toad rush	3.5 – 5.2 kg/ha	Apply a commercial knock down treatment one month prior to applying WEAPON so as to apply to bare ground.  Apply WEAPON within 1 month prior to planting for optimum weed control.  Use the higher rate for longer residual control.  <sup>1</sup> Kikuyu grass: use the higher rate for control for 3 months after application. <sup>2</sup> Red sorrel: use the higher rate for control up to 5 months after application.
	Post-planting	African daisy Common Storksbill Chickweed Fleabane Phalaris Silver grass Soft brome Spear thistle Toad rush		Apply WEAPON within 3 months of planting.  Use Uptake oil or another suitable wetter.
Eucalyptus	Pre-Planting	African daisy Common Storksbill Chickweed Fleabane Phalaris Silver grass Soft brome Spear thistle Toad rush	3.5 – 5.2 kg/ha	Apply a commercial knock down treatment one month prior to applying WEAPON so as to apply to bare ground.  Apply WEAPON within 1 month prior to planting for optimum weed control.
	Post-Planting	African daisy Common Storksbill Chickweed Fleabane <sup>3</sup> Phalaris Silver grass Soft brome Spear thistle Toad rush Witchgrass		3.5 kg/ha

### Additional Weeds List:

Burr medic,  
Black bindweed,  
Corn gromwell (White ironweed, Sheepweed),  
Deadnettle,  
Indian hedge mustard,  
Long haired poppy,  
Mint weed,  
Ox tongue,

Prickly lettuce,  
Shepherd's purse,  
Sow thistle,  
Turnip weed,  
Wild gooseberry,  
Wild radish,  
Wild turnip,  
Wireweed/hogweed.

[www.farmalinx.com](http://www.farmalinx.com)

CONTINUED OVERLEAF

containers or product.  
State or Territory government regulation. Do not burn empty  
vegetation and tree roots, in compliance with relevant Local  
and set up for this purpose clear of waterways, desirable  
500 mm below the surface in a disposal pit specifically marked  
management facility is not available bury the empty packaging  
to an approved waste management facility. If an approved waste  
recycling, break, crush or puncture and deliver empty packaging  
undiluted chemicals on-site. If recycling, replace cap and return  
clean containers to recycler or designated collection point. If not  
Store in the closed, original container in a dry, cool, well-  
ventilated area out of direct sunlight. Triple-rinse containers  
before disposal. Add rinsings to the spray tank. Do not dispose of  
PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET  
PLANTS  
DO NOT apply or drain or flush equipment on or near desirable  
trees or other plants or on areas where their roots may extend  
or in locations where the chemical may be washed or moved  
into contact with their roots. DO NOT apply on sites where  
contaminated clothing, and respirator and if rubber, wash with  
detergent and warm water.  
Wash hands after use. After each day's use, wash gloves,  
contaminated clothing, and respirator and if rubber, wash with  
detergent and warm water.  
at least 6 months following the last application.  
Store in the closed, original container in a dry, cool, well-  
ventilated area out of direct sunlight. Triple-rinse containers  
before disposal. Add rinsings to the spray tank. Do not dispose of  
undiluted chemicals on-site. If recycling, replace cap and return  
clean containers to recycler or designated collection point. If not  
recycling, break, crush or puncture and deliver empty packaging  
to an approved waste management facility. If an approved waste  
management facility is not available bury the empty packaging  
500 mm below the surface in a disposal pit specifically marked  
and set up for this purpose clear of waterways, desirable  
vegetation and tree roots, in compliance with relevant Local  
State or Territory government regulation. Do not burn empty  
containers or product.

### CONDITIONS OF SALE:

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Farmalinx Pty Ltd or under abnormal conditions. Farmalinx Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

## NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

### WITHHOLDING PERIOD

**Grazing: DO NOT graze any treated plants or cut for stock food**

### GENERAL INSTRUCTIONS

Farmalinx Weapon Herbicide may be applied pre-planting, preferably to bare soil, or post-planting of forestry seedlings to provide residual control of weeds to assist the establishment of the seedlings. Heavy rainfall on light soils may cause movement of the herbicide out of the weed seed zone, resulting in reduced weed control. WEAPON is primarily incorporated by rainfall. Sufficient rainfall (20 to 30 mm either as rainfall or irrigation) to wet the soil through the weed root zone is necessary within 2 to 3 weeks of application to make the product effective.

### RESISTANT WEEDS WARNING

FARMALINX WEAPON Herbicide is a member of the triazines group of herbicides. FARMALINX WEAPON has the inhibitors of photosynthesis at photostem II mode of action. For weed resistance management, WEAPON is a Group C herbicide. Some naturally-occurring weed biotypes resistant to WEAPON and other Group C herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by WEAPON or other Group C herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Farmalinx Pty Ltd accepts no liability for any losses that may result from the failure of WEAPON to control resistant weeds.

### MIXING

DO NOT mix or load within 20 m of any well, dam, intermittent or perennial stream. Partially fill the spray tank with water and maintain agitation while adding the required quantity of WEAPON. Agitation should continue after filling and during spraying. If spray is allowed to stand, agitate thoroughly before resuming spraying.

COMPATIBILITY  
This product is compatible with clopyralid and glyphosate (BAZOOKA). It is advisable that a small scale compatibility test be carried out prior to tank mixing with other products.

APPLICATION  
For aerial use, separate persons must perform the operations of mixing/loading and application.  
For aerial use, separate persons must perform the operations of mixing/loading and application.  
Apply with closed cockpit aircraft or closed cab tractor only.

GROUND APPLICATION  
Apply by ground boom in a minimum spray volume of 100 L/ha.  
Apply with closed cab tractor only.

ENVIRONMENT  
Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with the product or used containers.

PRE-PLANTING  
A commercial knockdown treatment should be applied 1 month prior to the application of WEAPON. Apply to moist soil or when rain is imminent and prior to germination of weeds up to a week before planting.

**farmalinx**  
Partners in Agriculture

FARMALINX Pty Ltd ABN 95 134 353 245  
Level 25, Suite 2506, Tower 2,  
101 Grafton Street, Bondi Junction NSW 2022  
T: 02 9389 2455 F: 02 9389 2844  
Website: [www.farmalinx.com](http://www.farmalinx.com)

APVMA Approval No. 88367/121065

**POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**WEAPON**  
FARMALINX  
HERBICIDE

ACTIVE CONSTITUENT: 875 g/kg TERBUTHYLAZINE

**GROUP C HERBICIDE**

For the control of weeds in forestry plantations as specified in the Directions for Use table.

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.

FARMALINX Pty Ltd ABN 95 134 353 245  
Level 25, Suite 2506, Tower 2, 101 Grafton Street,  
Bondi Junction NSW 2022

**farmalinx**