

**DIRECTIONS FOR USE****Restraint**

DO NOT treat physically damaged seed.

CROP	DISEASE/PEST	RATE	CRITICAL COMMENTS
Wheat	Bunt, Flag Smut (seed and soil borne), Loose Smut	100 mL/100 kg seed	Seed Dressing should be mixed with water to give even coverage of seed. The quantity of water will vary depending on type of equipment and quality of seed. Use a minimum of 400 mL of mixture (ie. Seed Dressing plus water) with each 100 kg of seed. DO NOT use more than 600 mL of mixture (product + water) with each 100 kg of seed. Whatever dilution is used it is essential that 100 mL of Seed Dressing is applied per 100 kg seed. The mixture should be gently stirred regularly. If nasal irritation occurs wear a face mask while treating seed.
Barley	Covered and Loose Smut		
Oats	Covered and Loose Smut		
Wheat, Barley, Oats	Protection against pests of stored grain: Granary Weevil, Lesser Grain Borer, Rice Weevil, Rust-red Flour Beetle, Sawtoothed Grain Beetle		

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

FOOD PRODUCING CROPS: NOT REQUIRED WHEN USED AS DIRECTED.

STOCK: DO NOT GRAZE PLANTS GROWN FROM TREATED SEED, OR CUT FOR STOCK FOOD WITHIN 4 WEEKS OF SOWING.

GENERAL INSTRUCTIONS

In Wheat: SPECTACLE may delay emergence. Crop will recover rapidly in moist soil types. However, treated seed should not be sown in soils that are likely to clod or crust over after sowing. SPECTACLE treated wheat deeper than 50 mm may slightly reduce percentage plant establishment in heavier soils and sowing rates may need to be adjusted accordingly. Germination may be reduced if poor quality seed is used (cracked or shrivelled) – best to use high quality seed. Treated seed may be held over for sowing in the following season. If so, additional insecticide treatment may be necessary. If the seed is not used immediately after treatment, it should be stored in a dry, well-ventilated place. Treated seed which will not be used for sowing at all should be destroyed.

FUNGICIDE RESISTANCE WARNING

Farmalinx SPECTACLE Seed Dressing is a Group 3 Fungicide and a member of the D.M.I. group of fungicides. For fungicide resistance management SPECTACLE is a Group 3 fungicide. Some naturally occurring individual fungi resistant to SPECTACLE and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by SPECTACLE or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Farmalinx Pty Ltd accepts no liability for any losses that may result from the failure of SPECTACLE to control resistant fungi.

GROUP 3 FUNGICIDE

INSECTICIDE RESISTANCE WARNING

For insecticide resistance management Farmalinx SPECTACLE Seed Dressing is a Group 15 Insecticide. Some naturally occurring insect biotypes resistant to SPECTACLE and other group 15 insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if SPECTACLE or other Group 15 insecticides are used repeatedly. The effectiveness of SPECTACLE on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Farmalinx Pty Ltd accepts no liability for any losses that may result from the failure of SPECTACLE to control resistant insects. SPECTACLE may be subject to specific resistance management strategies. For further information contact your local supplier, Farmalinx Pty Ltd representative or local agricultural department agronomist.

GROUP 15 INSECTICIDE

MIXING for use in a pickle pot or similar (liquid application)

The required quantity of SPECTACLE should be thoroughly mixed with a small quantity of clean, filtered water to make a creamy mix. Add this blend to the required quantity of water in the tank and mix thoroughly. Agitate prior to and during spraying.

DO NOT mix more than can be used for the day's seeding.

DO NOT tank mix with trace elements or other chemicals without first check compatibility.

SEED QUALITY

SPECTACLE should not be used on seed with more than 12% moisture content, as the treatment can raise the level by 0.6%. Seed to be treated should be of the highest quality. Poor quality (sprouted, cracked, pinched etc.) seed should not be used.

CLEANING OF EQUIPMENT

Clean equipment (augers etc.) with water after use. If changing from this water based product to a solvent based formulated product or vice versa, ensure complete dryness with Methylated spirits (or similar). Failure to do so will result in clogging of equipment.

PRECAUTION

DO NOT allow seed treated with this product to contaminate grain intended for human consumption. If possible, use a separate auger for treating seed. If using the same auger for treating grain and moving grain for human consumption, ensure auger is completely cleaned as above.

PROTECTION OF LIVESTOCK

Seed treated with this product must not be used for animal consumption, poultry feed or mixed with animal feed. DO NOT allow seed treated with this product to contaminate grain intended for animal consumption.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT feed treated seed or otherwise expose to wild or domestic birds or other animals. Harmful to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers. Used containers or bags which have held treated seed must be correctly disposed of.

STORAGE AND DISPOSAL

When treated seed is stored it should be kept apart from other grain. Such bags or other containers should clearly be marked to indicate the contents have been treated with SPECTACLE. DO NOT use seed treated with SPECTACLE for human consumption. Bags which have held seed treated with this product are not to be used for any other purpose.

For Non-Refillable Containers: Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

For Refillable Containers

Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs Farmalinx Pty Ltd should be advised immediately. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Wash hands after use.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131 126, New Zealand 0800 764 766.

SDS

Additional information is listed in the safety data sheet (SDS). A safety data sheet for Farmalinx SPECTACLE Seed Dressing is available from Farmalinx Pty Ltd on request. Call Customer Service on 02 9389 2455.

CONDITIONS OF SALE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Farmalinx Pty Ltd or under abnormal conditions. Farmalinx Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FARMALINX
Spectacle

SEED DRESSING

ACTIVE CONSTITUENTS: 25 g/L TEBUCONAZOLE
4 g/L TRIFLUMURON

GROUP 3 FUNGICIDE

Wheat: For control of bunt, flag smut and loose smut.
Barley: For control of covered and loose smut of barley.

Oats: For control of covered and loose smut.
For protection against insect pests of stored grain.

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.

FARMALINX Pty Ltd ABN 95 134 353 245
Level 25, Suite 2506, Tower 2, 101 Grafton Street,
Bondi Junction NSW 2022

farmalinx

farmalinx
Partners in Agriculture

FARMALINX Pty Ltd ABN 95 134 353 245

Level 25, Suite 2506, Tower 2,

101 Grafton Street, Bondi Junction NSW 2022

T: 02 9389 2455 F: 02 9389 2844

Website: www.farmalinx.com

APVMA Approval NO.: 89073/123543