FLAT SIZE: 332 MM WIDE X 448 MM DEEP



DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply in southeast Queensland unless in July or August. DO NOT apply in Mary Burnett in April, May, October or November. DO NOT apply by vertical sprayer.

DO NOT apply by fixed wing aircraft.

- DO NOT apply if heavy rains or storms are forecast within 3 days. DO NOT irrigate to the point of runoff for at least 3 days after application.

DO NOT apply to waterlogged soil.

DO NOT apply more than 205 kg Farmalinx Weapon Herbicide (182 kg terbuthylazine) per day by air, or 78 kg Farmalinx Weapon Herbicide (68.25 kg terbuthylazine) per day by ground boom.

Sprav Drift Restraints:

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift.

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone table/s below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application. DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DO NOT apply by a boom sprayer unless the following requirements are met:

Spray droplets are not smaller than a COARSE spray droplet size category.

• Minimum distances between the application site and downwind sensitive areas are observed (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for boom sprayers').

Buffer zones for boom sprayers

| Application rate | Boom height above the target canopy | Mandatory downwind buffer zones | | | |
|------------------|-------------------------------------|---------------------------------|-----------------------|------------------|--|
| | | Bystander areas | Natural aquatic areas | Vegetation areas | |
| Up to 5.2 kg/ha | 0.5 m or lower | 50 metres | 160 metres | 70 metres | |
| 3.5 kg/ha | 0.5 m or lower | 50 metres | 100 metres | 50 metres | |
| | 1.0 m or lower | 50 metres | 375 metres | 140 metres | |

DO NOT apply by aircraft unless the following requirements are met

· For application by helicopter only, not for application by fixed-wing aircraft.

Spray droplets are not smaller than a COARSE spray droplet size category.

 For maximum release height above the target canopy of 3 metres or 25 per cent of wingspan or 25 per cent of rotor diameter, whichever is the greatest, minimum distances between the application site and downwind sensitive areas are observed (see the following table titled 'Buffer zones for aircraft')

Buffer zones for aircraft

| Type of aircraft | Mandatory downwind buffer zones | | | |
|------------------|---------------------------------|-----------------------|------------------|--|
| | Bystander Areas | Natural aquatic areas | Vegetation areas | |
| Helicopter | 130 metres | 800 metres | 400 metres | |

| CROP | SITUATION | WEEDS | RATE | CRITICAL COMMENTS |
|---------------|---------------|--|-----------------|---|
| Pinus radiata | Pre-planting | African daisy Common Storksbill Chickweed Fleabane Kikuyu grass ¹ Phalaris Red sorrel ² Silver grass Soft brome Spear thistle Toad rush See Additional Weeds List below | 3.5 – 5.2 kg/ha | Apply a commercial knock down treatment one month prior to applying WEAPON so as to apply to bare ground. Apply WEAPON within 1 month prior to planting for optimum weed control. Use the higher rate for longer residual control. ¹ Kikuyu grass: use the higher rate for control for 3 months after application. ² Red sorrel: use the higher rate for control up to 5 months after application. |
| | Post-planting | African daisy Common Storksbill Chickweed Fleabane Phalaris Silver grass Soft brome Spear thistle Toad rush See Additional Weeds List below | | Use TechDeck or another suitable wetter. |
| Eucalyptus | Pre-Planting | African daisy Common Storksbill Chickweed Fleabane Phalaris Silver grass Soft brome Spear thistle Toad rush See Additional Weeds List below | 3.5 – 5.2 kg/ha | Apply a commercial knock down treatment one month prior to applying WEAPON so as to apply to bare ground. Apply WEAPON within 1 month prior to planting for optimum weed control. |
| | Post-Planting | African daisy Common Storksbill Chickweed Fleabane ³ Phalaris Silver grass Soft brome Spear thistle Toad rush Witchgrass See Additional Weeds List below | 3.5 kg/ha | Use TechDeck or another suitable wetter. NOTE: Some early crop phytotoxicity may be ob- served particularly on light soils. Heavy, intense rain- fall following application may cause crop damage. Avoid overlapping sprays and spraying-out corners. ³ Where fleabane may be a problem in the summer months apply in combination with clopyralid 750 g/ kg at 200 g/ha. |

Additional Weeds List:

| Burr medic, |
|--|
| Black bindweed, |
| Corn gromwell (White ironweed, Sheepweed), |
| Deadnettle, |
| Indian hedge mustard, |
| Long haired poppy, |
| Mint weed, |
| Ox tongue, |
| - |

Prickly lettuce, Shepherd's purse, Sow thistle. Turnip weed, Wild gooseberry, Wild radish, Wild turnip, Wireweed/hogweed.



CONTINUED OVERLEAF

betore planting.

rain is imminent and prior to germination of weeds up to a week

prior to the application of WEAPON. Apply to moist soil or when A commercial knockdown treatment should be applied 1 month

Pre - Planting

Apply in spray volume of 50 L/ha. Apply with closed cockpit aircraft only. mixing/loading and application. watercourses with the product or used containers. For aerial use, separate persons must perform the operations of Very toxic to aquatic life. DO NOT contaminate wetlands or ENVIRONMENT

Retrial Application .sd/J 001 to emulov spray wolume of 100 L/ha.

weeds. Clothing and gloves must be laundered after each day's for maintenance activities requiring direct contact with treated wrist and impervious footwear. Wear chemical resistant gloves

cotton overalls (or equivalent clothing) buttoned to the neck and

When entering areas where treated weeds are present, wear

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND

containers or product.

STNAJ9

State or Territory government regulation. Do not burn empty vegetation and tree roots, in compliance with relevant Local, and set up for this purpose clear of waterways, desirable 500 mm below the surface in a disposal pit specifically marked management facility is not available bury the empty packaging to an approved waste management facility. If an approved waste recycling, break, crush or puncture and deliver empty packaging clean containers to recycler or designated collection point. If not undiluted chemicals on-site. If recycling, replace cap and return before disposal. Add rinsings to the spray tank. Do not dispose of ventilated area out of direct sunlight. Triple -rinse containers Store in the closed, original container in a dry, cool, well-**JA2092ID DNA 3DA90T2**

at least 6 months following the last application. NOT plant crops other than those recommended on this label for containing, or to be planted with susceptible crops or plants. DU surface water from heavy rain can be expected to runoff to areas

into contact with their roots. DO NOT apply on sites where

or in locations where the chemical may be washed or moved

trees or other plants or on areas where their roots may extend

DO NOT apply or drain or flush equipment on or near desirable

0380 5422°

from FARMALIXY Pty Ltd on request. Call Customer Service on O2 safety data sheet for FAMALIAW WEAPON Herbicide is available A.(SDS) teens and the safety data sheet (SDS). A SOS

Centre. Phone Australia 131126, New Zealand 0800 764 766. If poisoning occurs, contact a doctor or Poisons Information **DIA TZRIA**

detergent and warm water.

For aerial use, separate persons must perform the ΝΟΙΤΑϽΙΔη

closed cockpit aircraft or closed cab tractor only. operations of mixing/loading and application. Apply with

Apply with closed cab tractor only.

Ground Application

be carried out prior to tank mixing with other products. (BAZOOKA). It is advisable that a small scale compatibility test This product is compatible with clopyralid and glyphosate

YTIJIBITA9M00

suitable wetter at label rates when weeds have emerged. post - Planting

WEAPON can be applied post-planting. Add Techdeck or other

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS **AUTHORISED UNDER APPROPRIATE LEGISLATION**

ВЕ-ЕИТВҮ РЕВІОD

WITHHOLDING PERIOD

Grazing: DO NOT graze any treated plants or cut for stock food

GENERAL INSTRUCTIONS

Farmalinx Weapon Herbicide may be applied pre-planting, preferably to bare soil, or post-planting of forestry seedlings to provide residual control of weeds to assist the establishment of the seedlings. Heavy rainfall on light soils may cause movement of the herbicide out of the weed seed zone, resulting in reduced weed control. WEAPON is primarily incorporated by rainfall. Sufficient rainfall (20 to 30 mm either as rainfall or irrigation) to wet the soil through the weed root zone is necessary within 2 to 3 weeks of application to make the product effective.

RESISTANT WEEDS WARNING

FARMALINX WEAPON Herbicide is a member of the triazines group of herbicides. FARMALINX WEAPON has the inhibitors of photosynthesis at photostem II mode of action. For weed resistance management,

GROUP 5 HERBICIDE

WEAPON is a Group 5 herbicide. Some naturally-occurring weed biotypes resistant to WEAPON and other Group 5 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by WEAPON or other Group 5 herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Farmalinx Pty Ltd accepts no liability for any losses that may result from the failure of WEAPON to control resistant weeds.

MIXING

DO NOT mix or load within 20 m of any well, dam, intermittent or perennial stream. Partially fill the spray tank with water and maintain agitation while adding the required quantity of WEAPON. Agitation should continue after filling and during spraying. If spray is allowed to stand, agitate thoroughly before resuming spraying.

SNOITCORE SAFETY DIRECTIONS

contaminated clothing, and respirator and if rubber, wash with Wash hands after use. After each day's use, wash gloves, and full facepiece respirator with dust cartridge or canister. to the neck and wrist, elbow-length chemical resistant gloves loading wear cotton overalls, over normal clothing, buttoned May irritate the eyes. Avoid contact with eyes. When mixing and

POISON

Seller warrants that the product conforms to its chemical

description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal

conditions of use. No warranty (other than non-excludable

statutory warranties) of merchantability or fitness for a particular

purpose, express or implied, extends to the use of product

contrary to label instructions, or under off-label permits not

endorsed by Farmalinx Pty Ltd or under abnormal conditions.

Farmalinx Pty Ltd accepts no liability for any loss or damage

farmalinx

Partners in Agriculture

FARMALINX Pty Ltd ABN 95 134 353 245

Level 25, Suite 2506, Tower 2, 101 Grafton Street, Bondi Junction NSW 2022

T: 02 9389 2455 F: 02 9389 2844

Website: www.farmalinx.com

APVMA Approval No. 88367/135967

arising from incorrect storage, handling or use.

CONDITIONS OF SALE:

