

Battleship

HERBICIDE



DIRECTIONS FOR USE: RESTRAINTS:

DO NOT apply to crops or weeds which are not actively growing or to plants which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (water-logged or drought affected) or previous herbicide treatment, as crop damage or reduced levels of control may result. DO NOT use in high winds. DO NOT spray if rain is likely to occur within four hours. DO NOT apply close to, or on areas, containing roots of desirable vegetation, where treated soil may be washed into areas growing, or to be planted to, desirable plants, or on sites where surface water from heavy rain can be expected to run off to areas containing, or to be planted to, susceptible crops or plants. DO NOT move soil which may have been sprayed to areas where desirable plants are to be grown.

Table 1: NEW SOUTH WALES and QUEENSLAND only

CROP	CROP GROWTH STAGE	WEEDS CONTROLLED	WEED GROWTH STAGE	RATE /ha	CRITICAL COMMENTS
Barley, Canary grass, Oats, Triticale, Wheat	Apply from early tillering (when the main shoot has 4-5 leaves and 2 or more new tillers have formed) to start of jointing (first node (Z30))	Climbing Buckwheat (Black Bindweed)	Up to 6 leaf	1L	For best control of Climbing Buckwheat apply at early tillering as this weed becomes increasingly difficult to control as it becomes larger. To avoid possible crop damage, DO NOT spray outside the early tillering to early jointing stage.
		Common Sowthistle Doublegee (Spiny Emex) Mustard Saffron Thistle Skeleton Weed Turnip Weed Variegated Thistle Wild Radish Wild Turnip	Young rosette or seedling plants up to 8 true leaves		
		NZ Spinach ¹ Wireweed ¹	Up to 4 leaf	¹ Suppression only - Spray early.	
Barley, Triticale, Wheat		Common Sowthistle NZ Spinach Wireweed	Apply up to 4 leaf stage	800 mL + 5 g Metfuron(600 g/kg metsulfuron methyl)	Add a Wetdrop wetter at the rate of 100 mL/100 L water.
Linseed Linola	Apply when crop is 8-20 cm high	Climbing Buckwheat (black bindweed) Doublegee (Spiny Emex) Mustard Saffron Thistle Skeleton Weed Variegated Thistle Wild Radish Wild Turnip	Young rosette or seedling plants up to 4 true leaves	670 to 840 mL	Use the higher rate when Climbing Buckwheat is present or weed growth is advanced.
		NZ Spinach ¹ Wireweed ¹	Up to 2 leaf		

Table 2: WESTERN AUSTRALIA only

CROP	CROP GROWTH STAGE	WEEDS CONTROLLED	WEED GROWTH STAGE	RATE /ha	CRITICAL COMMENTS
Barley, Canary grass, Oats, Triticale, Wheat	Apply from 3 leaf to early tillering (when the main shoot has 4-5 leaves and 2 or more new tillers have formed) to start of jointing (first node (Z30))	Capeweed Doublegee Saffron Thistle Turnip Weed Wild Mustard Wild Turnip	Young rosette or seedling plants up to 4 true leaves	600 mL	To avoid possible crop damage, DO NOT spray outside the early tillering to early jointing stage. Caution: DO NOT use this product where the land to be treated will be sown to lupins or peas the following year or where a volunteer medic or subclover pasture is required in the following season. Only use when a cereal crop will follow the cereal crop to be treated.
		Capeweed Doublegee Saffron Thistle Turnip Weed Wild Mustard Wild Turnip	Young rosette or seedling plants up to 6 true leaves	750 mL	
		Wireweed ¹	Up to 4 leaf		
	Apply from early to mid-tillering	Docks Sorrel	Active growth	1 L	¹ Suppression only - Spray early.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHOLDING PERIODS: DO NOT GRAZE OR CUT CROPS OR PASTURES FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION. HARVEST: NOT REQUIRED WHEN USED AS DIRECTED. MINIMUM RECROPPING PERIODS: PICLORAM, ONE OF THE ACTIVE CONSTITUENTS IN THIS PRODUCT, REMAINS ACTIVE IN THE SOIL FOR EXTENDED PERIODS DEPENDING ON RATE OF APPLICATION, SOIL TYPE, RAINFALL, TEMPERATURE, HUMIDITY, SOIL MOISTURE AND SOIL ORGANIC MATTER.

DO NOT PLANT SUSCEPTIBLE BROADLEAF CROPS WITHIN 12 MONTHS OF APPLYING THE PRODUCT. CEREAL CROPS AND GRASSES CAN BE SOWN SAFELY AFTER USING FARMALINX BATTLESHIP HERBICIDE-242 FOR IN CROP WEED CONTROL.

GENERAL INSTRUCTIONS

RESISTANT WEEDS WARNING Farmalinx Battleship Herbicide contains members of the pyridine and phenoxy groups of herbicides. The product has the disrupters of plant cell growth mode of action. For weed resistance management the product is a Group I Herbicide. Some naturally occurring weed biotypes resistant to the product and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Farmalinx Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture, or local Farmalinx Pty Ltd representative.

MIXING Farmalinx Battleship Herbicide may be mixed with water only. Mix only sufficient chemical for each day's use and avoid storing mix. Half fill the spray tank with water and add the required quantity of Farmalinx Battleship Herbicide and complete filling. Agitate continuously to ensure thorough mixing before and during application.

Tank Mixtures: Wettable powder or dry flowable formulations (eg. water dispersable granules) should be added to the spray tank first, followed by suspension concentrates (flowables), water soluble salts and then emulsifiable concentrate formulations (Flagship™). Add spraying oils and surfactants (wettors) last, if required.

NOT A DANGEROUS GOOD ACCORDING TO THE AUSTRALIAN DANGEROUS GOODS (ADG) CODE.

SMALL SPILL MANAGEMENT Wear protective equipment (See SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal. (See STORAGE AND DISPOSAL section).
NOTICE TO BUYER Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off-label permits not endorsed by FARMALINX Pty Ltd or under abnormal conditions. FARMALINX Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.
 *Other Registered Trademarks.

FIRST AID If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.
 Harmful if swallowed. Attacks eyes. Avoid contact with eyes.
SAFETY DIRECTIONS Harmful if swallowed. Attacks eyes. Avoid contact with eyes. Will irritate the skin. When preparing the spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and goggles. If product in eyes, wash it out immediately with water. After use wash thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.
 DO NOT burn empty containers or product.
STORAGE AND DISPOSAL Store in closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs, Farmalinx Pty Ltd should be advised immediately. This minibulk container is reusable and remains the property of Farmalinx Pty Ltd. DO NOT reuse empty container. Empty contents fully into application equipment. Close all valves and return to the point of supply for refill or storage. No other liquid, solid or pesticide product should be put into it. When empty return to Farmalinx Pty Ltd. for cleaning, relabelling and refilling.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT DO NOT contaminate streams, rivers or waterways with the chemical or used containers.
PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS DO NOT use on land to be cultivated for growing susceptible crops within 12 months of applying Farmalinx Battleship Herbicide. Legumes, vines, vegetables, cotton, tomatoes, ornamentals and many other plants are highly susceptible to this herbicide during both growing and dormant periods. Cereal crops and grasses can be sown safely after using Farmalinx Battleship Herbicide. This product will kill legumes (clovers, medics) present in the crop at the time of spraying. In the seasons following application of this product the regeneration or establishment of sensitive legumes (clover, medics, peas, lupins) may be adversely affected by soil residues. Cereal crops may be reduced in height following use of this product. This height reduction is a normal response and does not represent any yield loss.
 DO NOT allow spray drift onto susceptible crops, such as cotton, tomatoes, vines, fruit, potatoes, vegetables, ornamentals, tobacco, lupins and other legumes, safflower, sugar beet, hops, flowers or shade trees.
 DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Minimise spray drift by using low pressures and nozzles which DO NOT give a fine droplet size. DO NOT apply close to or on areas containing roots of desirable vegetation, where treated soil may be washed to areas growing, or to be planted to, desirable plants, or on sites where surface water from heavy rain can be expected to run off to areas containing or to be planted to, susceptible crops or plants. DO NOT move soil which may have been sprayed to areas where desirable plants are to be grown.

www.farmalinx.com

CONTINUED OVERLEAF

Partners in Agriculture

FARMALINX Pty Ltd ABN 95 134 353 245
 Level 25, Suite 2506, Tower 2,
 101 Grafton Street, Bondi Junction NSW 2022
 T: 02 9389 2455 F: 02 9389 2844
 Website: www.farmalinx.com

APVMA Approval No. 68004/57255

CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FARMALINX Battleship HERBICIDE

GROUP I HERBICIDE

ACTIVE CONSTITUENT: 420 g/L MCPA present as the potassium salt 26 g/L PICLORAM present as the potassium salt

For the control of climbing buckwheat, common sowthistle, skeleton weed, capeweed, doublegee and other broadleaf weeds in winter cereals and linseed crops, as specified in the Directions for Use table.

IMPORTANT: THIS LABEL IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.

FARMALINX Pty Ltd ABN 95 134 353 245
 Level 25, Suite 2506, Tower 2, 101 Grafton Street,
 Bondi Junction NSW 2022

farmalinx

COMPATIBILITY Farmalinx Battleship Herbicide is compatible with the following: (Read and follow all label directions, restraints, plant-back periods, withholding periods and safety directions on the partner label as well as those on the Farmalinx Battleship Herbicide label).

Broadleaf Herbicides
 Eclipse™ Herbicide; Clop 300 Herbicide; MCPA amine; MCPA LVE; Metfuron 600 WG; Floxor 200EC Herbicide; 2,4-D amine.

Grass Herbicides
 Diclofop-methyl (Annual Ryegrass only); Wildman 240EC (Wild Oats and Phalaris only)
 Adjuvants:- Wetdrop wetter/BS-1000 (when mixed with metsulfuron-methyl); Hotwire™ Spraying Oil (when mixed with Wildman 240 EC)

Insecticides
 Lemat™ ; Farmalinx Dimethoate

Farmalinx Battleship Herbicide is not compatible with zinc products (such as Zinc dextrase and Zinc heptahydrate), copper salts (such as sulphate) and manganese sulphate. DO NOT use with hard water or water clarified with alum.

APPLICATION METHODS
Ground application (Boom)
 Apply Farmalinx Battleship Herbicide with an accurately calibrated boom sprayer, in at least 50 L/ha water. Flat fan nozzles are recommended using pressures in the range 200 to 300 kPa. Set the boom at a height to ensure a double overlap of the nozzle pattern.

Aerial Application
 Apply in a minimum volume 35 L/ha water. Use equipment calibrated to produce spray droplets not smaller than COARSE according ASAE S572 Standard. DO NOT apply when the temperature is above 30°C, when there is no wind or when the wind is blowing toward susceptible crops.

Cleaning Spray Equipment:
 Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto wasteland away from desirable plants and water courses.

Rinsing:
 After using Farmalinx Battleship Herbicide, empty the tank completely and drain the whole system. Thoroughly wash inside the spray unit using a pressure hose. Drain, and clean any filters in the tank, pump, lines, hoses and nozzles. After cleaning the tank as above, quarter fill with clean water and circulate through the pump, lines and nozzles. Drain and repeat the rinsing procedure twice.

Decontamination (before spraying cotton and other sensitive crops; see PROTECTION OF CROPS): Wash the tank and rinse the system as above. Then quarter fill the tank and add an alkali detergent (eg. liquid Surf®, Omo® or Drive™) at 500 mL/100 L of water or the powder equivalent at 500 g/100 L and circulate throughout the system for at least 15 minutes. Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain.