FARMALINX Belta 700 WG

DIRECTIONS FOR USE RESTRAINTS:

DO NOT apply to very wet soils if rain is imminent or to soils prone to waterlogging. DO NOT apply to soils of very high organic matter content. DO NOT apply to crops or weeds under stress caused by factors such as root or foliar diseases, nutrient deficiencies, or extremes of temperature or moisture.

CROP	WEEDS CONTROLLED	STATE	RATE per ha	CRITICAL COMMENTS
Pre- emergence Chickpeas	Deadnettle (Lamium amplexicaule) Indian hedge mustard (Sisymbrium orientale) White ironweed (Buglossoides arvensis) Wild radish (Raphanus raphanistrum) Wireweed (Polygonum aviculare)		45 g plus 1.0 to 1.5 L Flowable Simazine Liquid (500 g/L)	Apply to moist, well prepared, clod and weed-free soil after planting and before crop emergence. Sufficient rainfall is required after application and prior to weed emergence to wet soil to a depth of 5 cm. Use the higher rate of simazine on heavier soils, or where higher weed pressure is expected, or where wireweed is a problem. Under adverse conditions, weeds may not be totally controlled but populations will be significantly reduced and surviving plants will generally be severely retarded. Good crop growth will aid weed control. Transient yellowing or reddening of the crop may occur. The risk of crop injury may be increased under adverse growing conditions. DO NOT use this mixture on soils, and in areas, ill suited to growing chickpeas, as crop injury will be increased.
Pre- emergence Faba beans Field peas	Indian hedge mustard (<i>Sisymbrium</i> <i>orientale</i>), Shepherd's purse (<i>Capsella bursa- pastoris</i>), Stinging nettle (<i>Urtica urens</i>) (NSW, Vic only), Toadrush (<i>Juncus</i> <i>bufonius</i>). Turrip weed (<i>Rapistrum</i> <i>rugosum</i>), * Deadnettle (<i>Lamium</i> <i>amplexicaule</i>), * Doublegee (<i>Emex</i> <i>australis</i>), * Paterson's curse (<i>Echium</i> <i>plantagineum</i>), * Wireweed (<i>Polygonum aviculare</i>), ** Wild oats (<i>Avena fatua</i>) * Wild radish (<i>Raphanus</i> <i>raphanistrum</i>), ** Annual ryegrass (<i>Lolium rigidum</i>), ** Barley grass (<i>Lolium rigidum</i>), ** Barley grass (<i>Hordeum leporinum</i>) Storksbill (<i>Erodium</i> spp.) Mouse-ear chickweed (<i>Carastium</i> <i>glomeratum</i>), * Capeweed (<i>Arctotheca alendula</i>), * Prickly lettuce (<i>Lactuca serriola</i>), * Yellow burrweed (<i>Amsinckia intermedia</i>) * Threehorn bedstraw (<i>Galium</i> <i>tricornutum</i>), * Wild radish (<i>Raphanus</i> <i>raphanistrum</i>), * Yellow burrweed (<i>Amsinckia intermedia</i>)	NSW, Vic, SA, WA only NSW, WA only SA, WA only NSW only SA only Vic, SA only	70 g	Apply to moist, well prepared, clod and weed-free soil after planting and before crop emergence. Sufficient rainfall is required after application and prior to weed emergence to wet soil to a depth of 5 cm. Under adverse conditions, weeds may not be totally controlled but populations will be significantly reduced and surviving plants will generally be severely retarded. Good crop growth will aid weed control. * Surviving plants will generally be retarded and will not compete with the crop. A follow-up spray with another product may be required for control of wild radish under high weed pressure or high rainfall conditions. ** Competition from grass weeds will be significantly reduced for at least 8 weeks. A post-emergence grass herbicide application may be required. (Refer to COMPATIBILITY section of label). *** Populations will not be reduced but plants will generally be significantly stunted. Seed set will also be reduced. Use the higher rates under anticipated high weed pressure and in high winter rainfall areas. DO NOT use on faba beans on light, sandy soils. On alkaline soils, the risk of crop damage to faba beans may be increased under adverse growth conditions. Refer to the CROP SAFETY and FOLLOW CROP sections of this label regarding varietal selectivity and follow crops.
	* Lesser loosestrife (<i>Lythrum</i> <i>hyssopifolia</i>), * Prickly lettuce (<i>Lactuca serriola</i>), ** Annual ryegrass (<i>Lolium rigidum</i>), ** Barley grass (<i>Hordeum leporinum</i>)		100 g	
Post- emergence Field peas (Following varieties only: Alma, Dun, Dundale, Early Dun, Wirrega)	Deadnettle (Lamium amplexicaule), Hare's ear (Conringia orientalis), Indian hedge mustard (Sisymbrium orientale), Toadrush (Juncus bufonius), * Wireweed (Polygonum aviculare)	NSW, Vic, SA only	70 g plus a non-ionic surfactant at 200 mL/ 100L	Apply to actively growing weeds in the cotyledon to 3-leaf stage. Weeds may not be totally controlled but populations will be significantly reduced and surviving plants will generally be severely retarded. Good crop growth will aid weed control. * Surviving plants will generally be retarded and will not compete with the area.
	* Threehorn bedstraw (Galium tricornutum)	Vic, SA only	70 to 100g plus a non- ionic surfactant at 200 mL/ 100L	the crop. Use the higher rate under high weed pressure. Refer to the FOLLOW CROP section of this label regarding follow crops.

CROP	WEEDS CONTROLLED	STATE	RATE per ha	CRITICAL COMMENTS
Pre- emergence Mungbeans, peanuts, soybeans	Common sida (Sida rhombifolia), Fat hen (Chenopodium album), Green amaranth (Amaranthus viridis), Pigweed (Portulaca oleracea), Redroot amaranth (Amaranthus retroflexus), Wild gooseberry (Physalis minima), * Anoda weed (Anoda cristata), * Belivine (Ipomoea plebeia), * Common thornapple (Datura stramonium), * Deadnettle (Lamium amplexicaule), * Jute (Corchorus olitorius), * Mintweed (Salvia reflexa), * Starburr (Acanthospermum hispidum), * Wild radish (Raphanus raphanistrum)	Qld, NSW, Vic, NT only	100 g	Apply to moist, well prepared, clod and weed-free soil after planting and before crop emergence. Sufficient rainfall or irrigation is required after application and prior to weed emergence to wet soil to a depth of 5 cm. Where soil crusting is likely, apply post-emergence. (Not mung beans). Under adverse conditions, weeds may not be totally controlled but populations will be significantly reduced and surviving plants will generally be severely retarded. Good crop growth will aid weed control. * Surviving plants will generally be retarded and will not compete with the crop.
	Above weeds plus Bladder ketmia (Hibiscus trionum), * Awnless barnyard grass (Echinochioa colona), * Apple of Peru (Nicandra physalodes), * Noogoora burr (Xanthium pungens), * Nutgrass (Cyperus rotundus)		140 g	
Post- emergence Peanuts, Soybeans	Common sida (Sida rhombifolia), wild gooseberry (Physalis minima), * Anoda weed (Anoda cristata), * Bathurst burr (Xanthium spinosum), * Bellvine (Ipomoea plebeia), * Common thornapple (Datura stramonium), * Deadnettle (Lamium amplexicaule), * Jute (Corchorus olitorius), * Fierce thornapple (Datura ferox)		100 g plus a non-ionic surfactant at 200 mL/ 100L	Apply to actively growing weeds in the 2 to 4 leaf stage. Weeds may not be totally controlled but populations will be significantly reduced and surviving plants will generally be severely retarded. Good crop growth will aid weed control. * Surviving plants will generally be retarded and will not compete with the crop. For grass weeds, a follow- up spray with a selective grass herbicide may be required.
	Above weeds plus Apple of Peru (Nicandra physalodes), Fat hen (Chenopodium album), * Awnless barnyard grass (Echinochloa colona), * Barnyard grass (Echinochloa crus- galli), * Nutgrass (Cyperus rotundus)		140 g plus a non-ionic surfactant at 200 mL/ 100L	
Pre- emergence to weeds Lucerne (established), serradella (established)	As for pre-emergence use in faba beans and field peas (winter weeds) and in soybeans (summer weeds)	Qld, NSW, Vic, SA, WA only	70 to 140 g	Use pre-emergence to weeds in established lucerne only . Apply following cutting or grazing, if necessary in mixtures with registered knock-down products. Apply at rates as per pre-emergence use in faba beans and field peas (winter weeds) and soybeans (summer weeds).
Centrosema (Cavalcade) (pre- emergence to crop)		NT only		Note CRITICAL COMMENTS applying to weed control in those crops.
Post- emergence Lucerne, serradella	As for post-emergence use in field peas (winter weeds) and in soybeans (summer weeds)	Qld, NSW, Vic, SA, WA only	70 to 140 g plus a non- ionic surfactant at 200 mL/ 100L	Apply to actively growing weeds in the cotyledon to 3 leaf stage (winter weeds) and 2-4 leaf stage (summer weeds). Apply at rates as per post- emergence use in field peas (winter weeds) and soybeans (summer weeds). Note CRITICAL COMMENTS applying to weed control in those crops. Seedling: Apply when crop is at the first trifoliate leaf stage or later (Autumn sown). Serradella varieties on which Farmalinx Belta 700 WG has been tested and found to be selective are: Avila, Elgara, Tauro.
Subterranean clover (Trifolium subterraneum)	Doublegee (Emex australis)	WA only	50 g plus 300 mL diuron (500 g/L) plus ammonium sulphate (417g/L) at 2 L per 100 L water	Established: Apply as above following cutting or grazing. Apply to actively growing weeds up to the 4 leaf stage and when the sub-clover is at the 3 leaf stage or later. This treatment should only be used in the first year of a pasture phase to aid sub-clover establishmen Weed numbers will be reduced and survivors will be stunted. Seed set will also be reduced. Sub-clover biomass may be reduced by this treatment. Varieties on which Farmalinx Belta 700 WG has been tested are Dalkeith and Nungarin. Other weeds may also be affected, thereby reducing the total pasture biomass.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIOD:

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION NOT REQUIRED WHEN USED AS DIRECTED GRAZING: HARVEST



CONTINUED OVERLEAF

higher than 100 g/ha were used in the previous crop.

+00 MOT plant these crops interim moisture (rainfall plus irrigation) from application to sowing is at least 800 mm. D0 NOT plant sorghum if Farmalianx Belta 700 WG rates DO NOT plant these crops unless interim moisture (rainfall plus irrigation) from application to sowing is at least 500 mm.

NOITADIJ99A AJTA ZHTNOM				
81	10	G	0	
all other crops (providing rainfall and	** maize (except for varieties with	chickpeas; lucerne; lupins; pasture	Maize varieties with CLEARFIELD	
(mm 000S sbeecks not inigation exceeds 2000 mm)		;səunbəj	Technology ONLY:	
	wnybjos **	* psrley;	- Pacific Hycorn 62IT	
		* wheat (except for wheat varieties with	- Pacific Hycorn 53IT	
		CLEARFIELD Technology);	- Pioneer 3395IR;	
		canola varieties with CLEARFIELD	mung beans; peanuts; soybeans	
		lechnology ONLY		

Following use in Summer crops: Irrigated only:

actually received, exclude single isolated heavy summer and autumn falls above 100 mm).

If expected rainfall is not received following use of Farmalinx Belta 700 WG, consult your local Farmalinx representative before planting wheat, barley or triticale. (In calculating rainfall

• In SAs and WB D0 V0T use on soils of PH 5.5 (Ca Cl₃) or less in areas where rainfall from spraying to sowing of cereals is expected to be below 400 mm. • In VSW, Vic and SB D0 V0T use the 100 g/ha rate in areas where rainfall from spraying to sowing of cereals is expected to be below 400 mm. mrsherm

* The following additional requirements apply if it is intended to sow WHEAT (except varieties with CLEARFIELD Technology), BARLEY or TRITICALE during the next winter season.
 * DO NOT apply Farmalinx Belta 700 WG pre-emergence later that the end of July.
 • DO NOT use Farmalinx Belta 700 WG in areas where rainfall from spraying to sowing of cereals is expected to be below 300 mm.

all other crops including canola (except rarieties with CLEARFIELD Technology; see 0 months)	oats; sattiower	lucerne; lupins; pasture legumes; (except varieties with CLEARFIELD (except varieties with CLEARFIELD	- Pacific Hycorn 62IT - Pacific Hycorn 53IT		
34	55	10	0		
MONTAE APPLICATION					

or damage arising from incorrect storage, handling or use.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off-label permits not sendorsed by FARMALINX Pty Ltd or under aborotable statutory varianties for the use of the product contrary to label instructions, or under off-label permits not sendorsed by FARMALINX Pty Ltd or under aborotable statutory warranties) or instructions of use use of the product contrary to label instructions, or under off-label permits not sendorsed by FARMALINX Pty Ltd or under aborotable statutory warranties) and the use of the product contrary to label instructions, or under off-label permits not sendorsed by FARMALINX Pty Ltd or under aborotable statutory warranties) and the use of the product contrary to label instructions, or under off-label permits not sendorsed by FARMALINX Pty Ltd or under aborotable statutory between the product contrary to label instructions, or under off-label permits not sendorsed by FARMALINX Pty Ltd or under aborotable statutory abortable statutory abortable statutory warranter abortable statutory abortable statutory abortable statutory abortable statutory warranter abortable statutory abortable statutory warranter abortable statutory abortable statutory abortable statutory abortable statutory abortable statutory abortable statutory warranter abortable statutory abort **CONDITIONS OF SALE**

For further information refer to the Material Safety Data Sheet (MSDS), which can be obtained from the supplier.

Sasm

If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 13 11 26.

DIA TZRIA

length PVC gloves and goggles. If product in eyes, wash it out immediately with water Wash hands after use. After each day's use, wash gloves, poggles and contaminated clothing. Will damage eyes. Will inritate skin. Avoid contact with the eyes and skin. When preparing spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow SAFETY DIRECTIONS

relevant Local, 5τατε or letritory government regulations. Do not burn empty containers or product.

If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging forom blow the sufficient of dispessal pit specifically marked and act up for this purpose clear of waterways, desirable vegetation and tree roots, in accordance with D0 N0T dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. original container in a dry, cool, well-yentilated area out of direct sunlight. Triple or preferably pressure rinse container before disposal. Add rinsings to spray tank. pesolo ent JA2092ID DNA 30A90T2

DO NOT contaminate dams, waterways or drains with this product or used containers.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

For the pre- or post-emergence control of certain weeds in Centrosema (Cavalcade), chickpeas, faba beans, field peas, lucerne, mung beans, peanuts, serradella, soybeans and subterranean clover as per the Directions For Use.

ACTIVE CONSTITUENT: 700 g/kg IMAZETHAPYR

HERBICIDE

m

GROUP

MPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER Gead Thoroughly before opening or using this product.

353 245 Street.

134 afton

ABN 95 1 2, 101 Gra

Ltd Towe 2022

X Pty | 2506, NSW 2

FARMALINX I Level 25, Suite 2: Bondi Junction M

armalinx

DO NOT plant sorghum if Farmalinx Belta 700 WG rates higher than 100 g/ha were used in the previous crop. *** D0 NOT plant these crops unless interim rainfall from application to sowing is at least 800 mm.

D0 N0T plant these crops unless interim rainfall from application to sowing is at least 500 mm.

 (\mathbf{r})

Following use in Winter crops:

The following minimum re-cropping intervals (months after application) should be observed.

Follow Crops: Under conditions, which do not favour breakdown of this product, roary-over soil residues can affect susceptible follow crops. As environmental and agronomic factors make it impossible to limitate all risks associated with the use of this product, rotational crop injury is always possible.

GENERAL INSTRUCTIONS

Farmalinx Belta 700 WG Herbicide can be used for either pre- or post- emergence weed control depending on the crop and weeds to be controlled. When the product is applied pre-emergence, susceptible weeds may emerge but growth will be retarded and weeds will either die or remain stunted and will not compete with the crop. Adequate soil moisture is important for optimum activity. When applied post-emergence, weeds will either die or remain stunted and will not compete with the crop.

A non-ionic surfactant, or an adjuvant as specified in the DIRECTIONS FOR USE, must be added to Farmalinx Belta 700 WG for post-emergence weed control. The addition of ammonium sulphate (417g/L) at a rate of 2 L/100 L water may assist in postemergence control of summer growing weeds.

MIXING

This product mixes readily with both hard and soft water. Fill the spray tank one half to three quarters full with clean water and then, with the agitator running, add the required amount of this product and then fill the tank with water. When tank mixing this product with other recommended compatible products, first add the other product to the tank and mix thoroughly before adding this product.

APPLICATION

Farmalinx Belta 700 WG should not be applied for a minimum of two hours before rainfall or irrigation

Ground Application: Avoid overlaps when spraying. Apply in 50 to 100 L/ha water using flat fan nozzles. The product may be applied in a band over the row in row crops. This will assist in minimising soil residue carry-over.

Aerial Application (Pre-emergence only): Avoid overlaps when spraying. Apply in a minimum 20 L/ha water in a maximum swath width of 18 m using a droplet VMD of 230-280 microns.

Do NOT apply Farmalinx Belta 700 WG under conditions (e.g. dead calm, excessive wind and/or small droplets) likely to cause spray drift onto wetlands or waterways, natural vegetation, crops other than specified on label or land to be planted with susceptible

EQUIPMENT CLEAN-UP

Thoroughly flush all spray equipment with water following use of Farmalinx Belta 700 WG and before use with other products.

Rinse water should NOT be discharged where it will reach streams, water-bodies or natural vegetation

COMPATIBILITY

This product is compatible with pendimethalin, oryzalin, alpha-cypermethrin, trifluralin, metribuzin, cyanazine, diuron, glyphosate, paraquat plus diquat, diquat, simazine, diflutenican, dimethoate, omethoate and endosulfan. D0 N0T tank mix with selective post-emergence grass herbicides. D0 N0T apply these herbicides following use of Farmalinx Belta 700 WG until grasses

have resumed active growth.

RESISTANT WEEDS WARNING Farmalinx Belta 700 WG Herbicide is a member of the Imidazolinone group of herbicides. Farmalinx Belta 700 WG has the inhibition of acetolactate synthase (ALS) mede of action. Farmalinx Belta 700 WG has the inhibition of acetolactate synthase (ALS) mode of action. For weed resistance management, Farmalinx Belta 700 WG is a Group B herbicides. Some naturally-occurring weed biotypes resistant to Farmalinx Belta 700 WG and other Group B herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population If these herbicides are used repeatedly. These weeds will not be controlled by Farmalinx Belta 700 WG or other Group B herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Farmalinx Pty Ltd accepts no liability for any losses that may result from failure of Farmalinx Belta 700 WG to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

D0 NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. D0 NOT spray within 50 m of wetlands or waterways.

Crop Safety: This product may cause slight shortening of plant internodes and may in some circumstances lead to transient crop yellowing but plants soon recover and yield is unaffected. This effect may be more pronounced when the product is used post-emergence or under poor growth conditions.

 $\rm DO\ NOT$ use this product pre-emergence on Collegian and Cressy Blue field pea

DO NOT use this product post-emergence on field pea varieties other than Alma, Dun, Dundale, Early Dun and Wirrega.

DO NOT use pre-emergence on serradella, seedling lucerne or subterranean clover.

DO NOT use post-emergence on chickpeas, faba beans or mung beans.

Should re-sowing of chickpeas, faba beans, field peas, lucerne, mung beans, peanuts, serradella, soybeans or subterranean clover be necessary, DO NOT re-apply Farmalinx Belta 700 WG.

NOITAJIAA RAPLICATION					
57	52	12	10	0	
all other crops (providing rainfall	cotton; maize (see also 0 months);	chickpeas; lucerne; lupins; pasture	** maize (except varieties with	Maize varieties with CLEARFIELD	
(mm 000S sbeeck				Technology ONLY:	
		* wheat (except for wheat varieties	wnybios **	- Pacific Hycorn 62IT - Pacific	
		with CLEARFIELD Technology);		Iycorn 53IT - Pioneer 3395IR; mung	
		canola varieties with CLEARFIELD		susest; peanuts; soybeans	
		Technology ONLY			

Dryland only: Dryland use the 140 gm rate in dryland soybeans, mung beans or peanuts unless it is intended to re-crop with a legumino CLEARFIELD Te

KEEP OUT OF REACH OF CHILDREN Read Safety Directions before opening or using

CAUTION

farmalinx

Partners in Agriculture

FARMALINX Pty Ltd ABN 95 134 353 245

Level 25, Suite 2506, Tower 2,

101 Grafton Street, Bondi Junction NSW 2022 T: 02 9389 2455 F: 02 9389 2844

> Website: www.farmalinx.com APVMA Approval No. 67670/56593

	רא המוני אין המוני אין המאמים אין המאמים אין המאמים אין המאמים אין אין האין אין האין אין האין אין האין אין האי					
[NOUTAFILGETER APPLICATION					
[22	55	12	10	0	
	all other crops (providing rainfall	; (sdfon; maize (see also 0 months);	chickpeas; lucerne; lupins; pasture	** maize (except varieties with	Maize varieties with CLEARFIELD	
	exceeds 2000 mm)			CLEARFIELD Technology);	lechnology ONLY:	
			* wheat (except for wheat varieties	wnyblos **	 Pacific Hycorn 62IT - Pacific 	
			(wnolondooT () I3I38A3 ID dtiwl		Innum 'Aldoss reennig - Tisa minni	