

FARMALINX

Bromoxy 200 EC

SELECTIVE HERBICIDE

DIRECTIONS FOR USE:

RESTRAINTS:
DO NOT apply by mister machines.
DO NOT apply if crops or weeds are stressed due to excessively dry or moist conditions.
DO NOT apply to diseased or frost-affected crops or if frosts are imminent.
DO NOT apply if rain is expected within three hours.
DO NOT apply to lucerne crops or clover based pastures or turf in temperatures above 20°C or when temperatures above 20°C may follow for some days after application.
DO NOT apply when windspeed is less than 3 or more than 20km per hour at application site.
DO NOT apply during surface temperature inversion conditions at the application site.
DO NOT apply with smaller than COARSE spray quality according to ASAE S572 definition for standard nozzles.

SITUATION & CROP	WEEDS CONTROLLED	WEED STAGE	STATE	RATE	CRITICAL COMMENTS
Wheat, Barley, Cereal Rye, Oats, Triticale, including those undersown with Medic (Vic only); Linseed, Clover or Lucerne Pastures; Lotus seed or forage crops (Old only)	Amsinckia (<i>Amsinckia</i> spp.), Black Bindweed (Climbing Buckwheat), (<i>Fallopia convolvulus</i>), Capeweed (<i>Arctotheca calendula</i>), Chamomile (<i>Matricaria matricarioides</i>), Common Peppercress (<i>Lepidium africanum</i>), Corn Gromwell (Ironweed), Sheepweed (<i>Buglossoides arvensis</i>), Fat Hen (<i>Chenopodium album</i>), Field Madder (<i>Sherardia arvensis</i>), Lesser Swinecress (<i>Coronopus didymus</i>), Mountain Sorrel (<i>Oxalis acetosella</i>), Purple Calandrinia (<i>Calandrinia menziesii</i>), Saffron Thistle (<i>Carthamus lanatus</i>), Shepherd's Purse (<i>Capsella burse-pastoris</i>), Three-horned Bedstraw (<i>Gallium tricornutum</i>), Tree Hogweed (<i>Polygonum patulum</i>), Variegated Thistle (<i>Silybum marianum</i>)	Up to the 4 leaf stage or when plants are no more than 35 mm in diameter	All States	1.4 L/ha	CROP STAGE: Wheat, Barley, Cereal Rye, Oats and Triticale: 3 leaf to full tillering (Z13-30). Linseed: 50 to 150 mm high. Lucerne & Clover: After seedlings have one trifoliolate leaf or before there is 150 mm regrowth in established stands. Lotus: Apply post-emergence at least 4 weeks after sowing (crop seedlings with a minimum of 2-6 true leaves). APPLICATION: Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application. Good coverage of the spray on the weed is essential. Apply only in the Autumn and Winter (mid April to end of August). For optimum control, apply only when the weeds are actively growing and before the weeds are shielded by the crop. Caution: Lucerne and Clover: This product will cause slight leaf burning of the undersown lucerne and clover. DO NOT apply to Persian Clover or Berseem Clover. Linseed: DO NOT spray Linseed by aerial application. Lotus: Some initial scorching of leaves may occur in some cases. Medics: Application to some cultivars of medics may result in unacceptable crop damage in some situations. # Effective control of Paterson's Curse and Fumitories will only be obtained under good growing conditions. ## Where Wireweed is growing under stress, it has been found to be less susceptible.
	Mexican Poppy (<i>Argemone ochroleuca</i>), Three Cornered Jack (Doublegee, Spiny Emex) (<i>Emex australis</i>), Turnip Weed (Giant Mustard) (<i>Rapistrum rugosum</i>), Wild Radish (<i>Raphanus raphanistrum</i>), Wild Turnip (<i>Brassica tournefortii</i>), Wireweed## (<i>Polygonum aviculare</i>), Paterson's Curse## (Salvation Jane) (<i>Echium plantagineum</i>)	Up to the 6 leaf stage or when plants are no more than 50 mm in diameter		2.0 L/ha	
	Fumitories# (<i>Fumaria</i> spp.)			Old, NSW, ACT, Vic, SA, WA only	
	Pheasant's Eye (<i>Adonis microcarpa</i>), Birds Eye (Carrot Weed) (<i>Cotula australis</i>)	Up to the 4 true leaf stage	Greater than the 4 true leaf stage	SA only	700 mL/ha
Fireweed (<i>Senecio</i> spp.)	Prior to flower development	Early flowering stage	Old, NSW, ACT only	1.4 L/ha	Apply during the Autumn-Winter period when the weeds are young and actively growing.
				2.8 L/ha	
Wheat, Barley, Cereal Rye, Oats, Triticale, including those undersown with Clover, Lucerne or Medics; Clover Seed Crops	Amsinckia (<i>Amsinckia</i> spp.), Capeweed (<i>Arctotheca calendula</i>), Chamomile (<i>Matricaria matricarioides</i>), Charlock (<i>Sinapis arvensis</i>), Corn Gromwell (Ironweed), Sheepweed (<i>Buglossoides arvensis</i>), Fat Hen (<i>Chenopodium album</i>), Field Madder (<i>Sherardia arvensis</i>), Hexham Scent (King Island Mellilot) (<i>Mellilotus indicus</i>), Horned Poppy (<i>Glaucium flavum</i>), Indian Hedge Mustard (<i>Sisymbrium orientale</i>), Lesser Swinecress (<i>Coronopus didymus</i>), Mexican Poppy (<i>Argemone ochroleuca</i>), Mintweed (<i>Salvia reflexa</i>), Purple Calandrinia (<i>Calandrinia menziesii</i>), Paterson's Curse (Salvation Jane) (<i>Echium plantagineum</i>) Rough Poppy (<i>Papaver hybridum</i>), Saffron Thistle (<i>Carthamus lanatus</i>), Shepherd's Purse (<i>Capsella burse-pastoris</i>), Slender Thistle (<i>Carduus tenuiflorus</i>), Turnip Weed (Giant Mustard), (<i>Rapistrum rugosum</i>), Wild Radish (<i>Raphanus raphanistrum</i>)	Up to the 5 leaf stage or when plants are no more than 50 mm in diameter	WA only	750 mL/ha PLUS 440 mL 2, 4-D Amine 625 per ha OR PLUS 700 mL MCPA 500/ha	CROP STAGE: Cereals: 3 leaf to full tillered (Z13-30). Clover: Apply after clover seedlings have at least three trifoliolate leaves. APPLICATION: Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application. Apply only in the Autumn and Winter (mid April to end of August). Apply only when the weeds are actively growing and before the weeds are shielded by the crop. Caution: Some damage and/or yield loss may occur in Olympic and Shortim wheat cultivars. DO NOT apply to Persian Clover or Berseem Clover. A slight burning of clover may occur. DO NOT use 2, 4-D or MCPA tank mix if wheat, oats, barley or triticale are undersown with lucerne, clover or medics unless some damage is acceptable. This product may be tank mixed with diclofop-methyl for control of Annual Rye Grass and Wild Oats in wheat and barley only. DO NOT use 2, 4-D or MCPA tank mix if using a tank mix with diclofop-methyl.
	Common Peppercress (<i>Lepidium africanum</i>), Tree Hogweed (<i>Polygonum patulum</i>)			Vic only	
	Black Bindweed (Climbing Buckwheat), (<i>Fallopia convolvulus</i>)	Up to the 8 leaf stage		Vic, SA only	

COMPATIBILITY
This product may be mixed in the spray tank with one of the following herbicides & insecticides in wheat and barley only: Diclofop Methyl Selective Herbicide, Dimethoate, Chlorpyrifos, Difenzoate, Dithiopyr, MCPA 500, Chlorsulfuron, Metsulfuron-Methyl, Diflufenican, Glyphosate, Propaquizafop and Dicamba.
PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS
DO NOT apply under weather conditions, or from spraying equipment, that may cause injury to crops, native and other non-target plants.
Note: Maximum daily temperatures at or above 20°C at application, or on days following treatment may result in seedling mortality and established plants may be damaged.
PROTECTION OF LIVESTOCK
Low hazard to bees. May be applied at any time as recommended in the Directions for Use.
PROTECTION OF WILDLIFE, FISH, CRUSTACEAN AND THE ENVIRONMENT
Dangers to users: DO NOT contaminate streams, rivers or waterways with the chemical or used containers.
STORAGE AND DISPOSAL
Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.
For Non-Refillable Containers: Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of unlabeled chemicals on site. If recycling, replace cap and return clean containers to recycler or a designated collection point. If not recycled, empties to be used for the purpose of a particular purpose. Expressions of opinion, extends to the use of the product contrary to label instructions, or under other labels, permits not endorsed by FARMALINX Pty Ltd or under abnormal conditions. FARMALINX Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.
* Other trademarks and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

GENERAL INSTRUCTIONS
Before opening, carefully read Directions for Use, Precautionary and Protection Statements, Storage & Disposal, Safety Directions and First Aid Instructions. Use only recommended growth stages of crops and weeds.
DO NOT use on crops other than recommended.
RESISTANT WEEDS WARNING
FARMALINX Bromoxy 200 EC Selective Herbicide is a member of the triazole group of herbicides. The product has the inhibitors of photosynthesis as its mode of action. For weed resistance management, the product is a Group C herbicide.
Some naturally occurring weed biotypes resistant to the product and other Group C herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by the product or other Group C herbicides.
Since the occurrence of resistance weeds is difficult to detect prior to use, FARMALINX Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds.
MIXING INFORMATION
To ensure even mixing, agitate half fill spray tank with clean water and add the required amount of this product. Agitate thoroughly, then add the remainder of the water. Agitate again before spraying commences.

BOOM SPRAYER
USE only low boom equipment set up to provide good coverage of weeds within the crop canopy.
USE ONLY coarse spray according to ASAE S572 definition for standard nozzles. Apply 50 to 200 (WA 50 to 100 litres) litres of water per hectare. Calibrate the machine before use to ensure that the correct application rate is obtained. Check height of boom above the crop to ensure that the spray is evenly distributed.
DO NOT use rain-drop or air induction nozzles as coverage of weeds may not be accurate.
AERIAL APPLICATION
Apply a minimum of 22 litres of water per hectare. Markers should always move upwind before aircraft passes overhead. Application by microaer equipment may give variable results.
EQUIPMENT MAINTENANCE
After use wash equipment thoroughly and rinse several times prior to re-use. Wash sprayer thoroughly with clean water after use. If using tank mixtures with 2,4-D Amine or MCPA 500 apply with equipment that will not be used in 2,4-D and MCPA susceptible crops, particularly cotton and vines.

FIRST AID
If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, do NOT induce vomiting. Give a glass of water.
SDS
Additional information is listed in the safety data sheet (SDS). A safety data sheet for FARMALINX Bromoxy 200 EC Selective Herbicide is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 9389 2455.
NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, or under other labels, extends to the use of the product contrary to label instructions, or under other labels, permits not endorsed by FARMALINX Pty Ltd or under abnormal conditions. FARMALINX Pty Ltd accepts no liability for any loss or damage arising from incorrect use, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield.
DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 8 WEEKS AFTER APPLICATION.
UNDER APPROPRIATE LEGISLATION, CONTRARY TO THIS LABEL UNLESS AUTHORISED

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Bromoxy 200 EC
FARMALINX

SELECTIVE HERBICIDE

200 g/L BROMOXYNIL present as the n-octanoylester
SOLVENT: 616 g/L LIQUID HYDROCARBON

GROUP C HERBICIDE

For the control of weeds in wheat, oats, barley, cereal rye and triticale (alone or undersown), linseed, clover and lucerne pastures; turf; fallow, non-crop, roadsides and rights-of-way as specified in the Directions for Use table.

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.

FARMALINX Pty Ltd ABN 95 134 353 245
Bond Junction NSW 2022

farmalinx

SITUATION & CROP	WEEDS CONTROLLED	WEED STAGE	STATE	RATE	CRITICAL COMMENTS	
Wheat, Oats, Barley, Triticale, alone or with undersown lucerne or clover	Common Sowthistle (<i>Sonchus oleraceus</i>)		Old, Northern NSW only		APPLICATION: Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application. Good coverage of the spray on the weed is essential. Apply only in the Autumn and Winter (mid April to end of August). For optimum control, apply only when the weeds are actively growing and before the weeds are shielded by the crop. Caution: Lucerne and Clover: This product will cause slight leaf burning of the undersown lucerne and clover. DO NOT apply to Persian Clover or Berseem Clover.	
		Up to 4 leaves		1.4 L/ha		Strictly for use on small weeds early in crop before shading out. CROP STAGE: Wheat, Barley, Oats and Triticale: 3 leaf to mid tillering (Z13-23). Use a minimum spray volume of 70 L/ha. If Group B resistance is suspected, or weeds have grown through a previous application of chlorsulfuron, apply 2.1 L/ha.
		4 to 8 leaves		2.1 L/ha		CROP STAGE: Wheat, Barley, Oats and Triticale: 3 leaf to full tillering (Z13-30). This product may cause slight leaf burning of undersown lucerne and clover.
Wheat, Barley, Oats and Triticale	Three Corned Jack (Doublegee, Spiny Emex) (<i>Emex australis</i>), Wireweed (<i>Polygonum aviculare</i>), Variegated Thistle (<i>Silybum marianum</i>)	Up to the 4 leaf stage or when plants are no more than 50 mm in diameter	Old, NSW, ACT, SA, Tas only	1.4 L/ha PLUS 440 mL 2, 4-D Amine 625 per ha OR PLUS 700 mL MCPA 500/ha	CROP STAGE: 5 leaf to full tillering (Z15-30). Apply under good growing conditions. BOOM SPRAYING: Use a minimum of 220 L spray per hectare on Black Bindweed when crop is greater than 300 mm high. Caution: Some crop damage and/or yield loss may occur in Olympic and Shortim wheat cultivars. DO NOT use 2, 4-D or MCPA tank mix if wheat, oats, barley or triticale are undersown with lucerne, clover or medics unless some damage is acceptable. DO NOT apply to medics (WA only). This product may be tank mixed with diclofop-methyl for control of Annual Rye Grass and Wild Oats in wheat and barley only. DO NOT use 2, 4-D or MCPA tank mix if using a tank mix with diclofop-methyl.	
	Fumitories (<i>Fumaria</i> spp.)					Old, NSW, ACT, Vic only
	Wild Mustard (<i>Sisymbrium</i> spp.), Wild Turnip (<i>Brassica tournefortii</i>)					Old, Vic only
	Cleavers	2-4 stems + 1-3 whorls or leaves/ stems	Old, NSW, ACT, Vic, Tas only	2.4 L/ha PLUS 440 mL 2, 4-D Amine 625 per ha OR PLUS 700 mL MCPA 500/ha		
	Three Corned Jack (Doublegee, Spiny Emex) (<i>Emex australis</i>), Wireweed (<i>Polygonum aviculare</i>)	4-5 leaf stage	WA only			
	Fumitories (<i>Fumaria</i> spp.)		Old, NSW, ACT, Vic, Tas, SA only			
Black Binweed (Climbing Buckwheat) Chamomile (<i>Matricaria matricarioides</i>), Fat Hen (<i>Chenopodium album</i>), Field Madder (<i>Sherardia arvensis</i>), Lesser Swinecress (<i>Coronopus didymus</i>), Mountain Sorrel (<i>Oxalis acetosella</i>), Purple Calandrinia (<i>Calandrinia menziesii</i>), Paterson's Curse (<i>Salvation Jane</i>) (<i>Echium plantagineum</i>), Shepherd's Purse (<i>Capsella burse-pastoris</i>), Three-horned Bedstraw (<i>Gallium tricornutum</i>), Variegated Thistle (<i>Silybum marianum</i>)	5-8 leaf stage	All States				

www.farmalinx.com

CONTINUED OVERLEAF

SITUATION & CROP	WEEDS CONTROLLED	WEED STAGE	STATE	RATE	CRITICAL COMMENTS
Fallow and Non Crop	Cowvine or Peachivine (<i>Ipomoea lonchophylla</i>)	Up to 6 leaves	Old, NSW, ACT only	1.4 L/ha to 2.1 L/ha	For all fallow and non-crop situations. Complete coverage is essential, plants which are shaded by stubble or other weeds will not be controlled. Weeds that are drought stressed and sprayed under hot conditions may not be adequately controlled. The addition of glyphosate will provide broader spectrum weed control where needed. Refer to the Compatibility section for appropriate glyphosate products. The rate of glyphosate used should be robust for the conditions and sufficient to control the hardest-to-kill weed in the situation. For general fallow weed control application volumes above 50 L/ha are recommended.
	Climbing Buckwheat (<i>Fallopia convolvulus</i>)	Cotyledon to mature, vining plants		1.5 L/ha alone or 1-1.5 L/ha plus glyphosate	
Pastures	Volunteer Conventional Cotton (<i>Gossypium hirsutum</i>)	Cotyledon to 6 leaves	Old, NSW, WA, NT only	1.5 L/ha alone or 1-1.5 L/ha plus glyphosate	For Volunteer Cotton control, including Roundup Ready varieties a minimum spray volume of 80 L/ha should be used. Bromoxy 200 EC alone or with glyphosate will not provide satisfactory control of ratoon or "slub" cotton.
	Volunteer Roundup Ready* Cotton (<i>Gossypium hirsutum</i>)	Cotyledon to 6 leaves		1.5 L/ha alone or plus glyphosate	
	Adonis (Pheasants Eye), Birds Eye (Carrot Weed)	Up to 4 leaves		SA only	
Pastures, Roadsides, Rights-of-way	Doublegee, Wild Radish	Up to 4 leaves	WA only	1.5 L/ha	Apply in May-June. Medics may be severely scorched at the low rate and killed at the high rate. DO NOT apply to polymorpha medics. Avoid application to clover based pastures in temperatures over 20°C or when temperatures above 20°C may follow for some days after application, as seedling mortality may occur and established plants may be damaged.
	Matricaria	Up to 3 leaves		750 mL/ha	
	Fireweed	Prior flower bud		Old, NSW, ACT only	
Pastures, Roadsides, Rights-of-way	African Daisy (<i>Senecio pterophorus</i>)	Up to and including the flowering stage	SA only	300 mL per 100 L water	The plants including the stems, must be thoroughly wet to the point of run-off. Apply when the plants are actively growing. Spraymate Activator should be added as 125 mL per 100 L of spray. A follow-up treatment may be necessary for large, well-established plants. Ensure overall coverage on weed seedling.
	Boneseed/Bitou-bush (<i>Chrysanthemoides molliflora</i>)	Seedlings only		Vic, Tas only	

CRITICAL COMMENTS
Apply evenly over the area to be treated. DO NOT mow turf for two days after treatment. A slight burning of grasses, especially Paspalum, may occur under hot conditions. knapsack 15 L 60 mL per 100 m² per 300 mL per/100 L water per ha

CRITICAL COMMENTS
Apply evenly over the area to be treated. DO NOT mow turf for two days after treatment. A slight burning of grasses, especially Paspalum, may occur under hot conditions. knapsack 15 L 60 mL per 100 m² per 300 mL per/100 L water per ha