

ClearView 200 GR

HERBICIDE

DIRECTIONS FOR USE

RESTRAINTS: DO NOT apply this product to salt or erosion prone areas.
DO NOT apply this product within 100 metres of a recognised water course.
DO NOT apply FARMALINX CLEARVIEW 200 GR Herbicide on land with a slope greater than 20 percent (11 degrees).
DO NOT apply FARMALINX CLEARVIEW 200 GR Herbicide under conditions which will cause pellet movement to non-target areas during application.
RETAIN at least 20 percent of the original tree population in wildlife corridors that are at least 100 metres wide.
 These are essential for shade and protection of livestock and native fauna. Ideally, shelter belts should traverse variable terrain, particularly ridgelines and hill tops as well as water courses, and link other vegetated corridors where possible.
FOR LONG TERM VIABILITY retain a minimum of 5 hectares for shade areas, regardless of vegetation type.
DO NOT apply FARMALINX CLEARVIEW 200 GR Herbicide on field crops, near desirable trees or shrubs or to areas into which their roots may extend or in locations where the chemical may be washed into contact with their roots as injury or death may occur.
DO NOT apply FARMALINX CLEARVIEW 200 GR Herbicide to land to be put under field crops within 5 years of application.

Table 1: Hand Application (WA and NT only)
See GENERAL INSTRUCTIONS – APPLICATION section for application method details.

SITUATION	WEEDS CONTROLLED	RATE g/m ²	CRITICAL COMMENTS
Tea tree regrowth and associated woody weeds on grazing land	Black Tea Tree	1.5	Estimate the area within 30cm beyond the drip zone of each target weed or group of weeds and calculate the amount of FARMALINX CLEARVIEW 200 GR Herbicide to cover area to be treated. Distribute the required dose uniformly within this area. Use the higher rate for heavy density regrowth and heavy clay soils. * Some suppression of these species is provided by the rates given.
	Broadleaf Tea Tree	1	
	Cocky Apple*	2	
	Paperbark Tea Tree, Poplar Gum	1.5	
	Swamp Box	2	
Other woody weeds on grazing land	African Boxthorn		
	Coolibah	1	
	Parkinsonia	1 to 1.5	
	Prickly Acacia, Rubbervine	1.5	
	Whitewood	1	
Giant sensitive tree on grazing land and other grasslands	<i>Mimosa pigra</i>	1	Apply after the onset of storms (to moisten soil, close soil cracks (where applicable) and freshen growth of giant sensitive tree) and before wet season surface flooding. Repeat application may be required in year two for long term control. Some grass suppression may occur.

Table 2: Hand Application (Qld only)
See GENERAL INSTRUCTIONS – APPLICATION section for application method details.

SITUATION	WEEDS CONTROLLED	RATE g/m ²	CRITICAL COMMENTS
Brigalow regrowth and associated problem woody weeds on grazing land	Belah	1.5	Estimate the area within 30cm beyond the drip zone of each target weed or group of weeds and calculate the amount of FARMALINX CLEARVIEW 200 GR Herbicide to cover area to be treated. Distribute the required dose uniformly within this area. Use the higher rate for heavy density regrowth and heavy clay soils. * Some suppression of these species is provided by the rates given.
	Brigalow, Currant Bush	1 to 1.5	
	Dawson Gum	1	
	False Sandalwood*	2	
	Holly Bush	1.5	

SITUATION	WEEDS CONTROLLED	RATE g/m ²	CRITICAL COMMENTS
Brigalow regrowth and associated problem woody weeds on grazing land	Limebush, Poplar Box, Whitewood, Yellowwood	1	Estimate the area within 30cm beyond the drip zone of each target weed or group of weeds and calculate the amount of FARMALINX CLEARVIEW 200 GR Herbicide to cover area to be treated. Distribute the required dose uniformly within this area. Use the higher rate for heavy density regrowth and heavy clay soils. * Some suppression of these species is provided by the rates given. Use only for regrowth control, not clearing of virgin bush.
	Tea tree regrowth and associated woody weeds on grazing land		
Tea tree regrowth and associated woody weeds on grazing land	Black Tea Tree	1.5	
	Broadleaf Tea Tree	1	
	Cocky Apple*	2	
	Paperbark Tea Tree, Poplar Gum	1.5	
	Swamp Box	2	
	Other woody weeds on grazing land		
Other woody weeds on grazing land	African Boxthorn	2	
	Silver Leaved Ironbark, Brown Box	1.5	
	Coolibah	1	
	Gidgee	0.5 to 1	
	Gum-topped Box	1.5	
	Groundsel Bush	1	
	Lantana*	2	
	Parkinsonia	1 to 1.5	
	Pink Bloodwood	1	
	Prickly Acacia, Rubbervine, Wild Rosemary	1.5	

Table 3: Hand Application (NSW only)
See GENERAL INSTRUCTIONS – APPLICATION section for application method details.

SITUATION	WEEDS CONTROLLED	RATE g/m ²	CRITICAL COMMENTS
Brigalow regrowth and associated problem woody weeds on grazing land	Brigalow, Currant Bush, Holly Bush, Limebush, Whitewood	1 to 1.5	Estimate the area within 30cm beyond the drip zone of each target weed or group of weeds and calculate the amount of FARMALINX CLEARVIEW 200 GR Herbicide to cover area to be treated. Distribute the required dose uniformly within this area. Use the higher rate for heavy density regrowth and heavy clay soils. * Some suppression of these species is provided by the rates given. ** DO NOT use within 30m of trees. DO NOT apply to single continuous area greater than 0.5 hectares in size.
	False Sandalwood*	2	
	Other woody weeds on grazing land		
	African Boxthorn		
	Blue Heliotrope**	0.5	
Other woody weeds on grazing land	Groundsel Bush	1	
	Lantana*	2	
	Wild Rosemary	1.5	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

FOR NATIVE VEGETATION
 Use of FARMALINX CLEARVIEW 200 GR Herbicide on native vegetation must be done in accordance with STATE and/or LOCAL legislation, including the Vegetation Management Act 2000 in Queensland.

WITHOLDING PERIODS: NOT REQUIRED WHEN USED AS DIRECTED.

can be minimised by application when grasses are dormant (autumn application). FARMALINX CLEARVIEW 200 GR Herbicide may seriously injure desirable forage plants such as pasture legumes and other broadleaf plants. These species may regenerate in subsequent seasons.

PROTECTION OF LIVESTOCK
 Poisonous plants may become more palatable after application of granules and stock should be kept away from these plants until they have died down.
PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT
 DO NOT contaminate streams, rivers or waterways with the chemical or herbicide.
STORAGE AND DISPOSAL
 Store in the closed original container in a dry, cool well-ventilated area out of direct sunlight.
FIRST AID
 If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.
MATERIAL SAFETY DATA SHEET
 Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FARMALINX CLEARVIEW 200 GR Herbicide is available from Farmalinx Pty Ltd on request. Call Customer Service on 02 9389 2455.

NOTICE TO BUYER
 Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off-label permits not endorsed by FARMALINX Pty Ltd or under abnormal conditions. FARMALINX Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

farmalinx
 Partners in Agriculture

FARMALINX Pty Ltd ABN 95 134 353 245
 Level 25, Suite 2506, Tower 2,
 101 Grafton Street, Bondi Junction NSW 2022
 T: 02 9389 2455 F: 02 9389 2844
 Website: www.farmalinx.com

APVMA Approval No. 69043/59704

POISON
 KEEP OUT OF REACH OF CHILDREN
 READ SAFETY DIRECTIONS BEFORE OPENING OR USING

ClearView 200 GR
 HERBICIDE

ACTIVE CONSTITUENT: 200 g/kg TEBUTHIURON

GROUP C HERBICIDE

For control of brigalow re-growth, tea tree re-growth, Mimosa pigra and certain problem woody weeds on grazing lands as specified in the Directions For Use table by hand application.

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.

FARMALINX Pty Ltd ABN 95 134 353 245
 Level 25, Suite 2506, Tower 2, 101 Grafton Street,
 Bondi Junction NSW 2022

farmalinx

GENERAL INSTRUCTIONS
 FARMALINX CLEARVIEW 200 GR Herbicide may be applied at any time of the year. Treatments become effective after sufficient rainfall has occurred to move the chemical into the root zone where it is absorbed by woody plants. Applications just prior to seasonal rainfall give the most rapid response.
 Herbicidal activity will continue for several seasons after application. During this time woody plants may go through repeated defoliations and regrowths until death occurs. A single application is normally effective for several years. Forage grass production usually increases as woody weed competition is reduced. Increased grass production is also dependant on adequate rainfall and an acceptable grazing management programme. It is important to allow grasses to seed and seedlings to become established after treatment before grazing pressure is increased. Areas treated with FARMALINX CLEARVIEW 200 GR Herbicide may be overseeded with suitable pasture species. Consult your agricultural specialist for details on suitable seeding rates, timing and fertiliser application programme.
 The residual action of FARMALINX CLEARVIEW 200 GR Herbicide may be impaired if fire occurs before clay pellets are dissolved and the chemical has moved into the root zone.


APPLICATION
 It is recommended that gloves be used during hand application. Apply FARMALINX CLEARVIEW 200 GR Herbicide by uniformly distributing by hand over the desired area after weighing out the recommended quantity of pellets. The best way to calibrate yourself is to weigh a normal handful of pellets. If, for example, a handful weighs 20 grams, then this will treat 20 to 40m² at the rate of 0.5 to 1g/m².


DISPENSER PACK
 Apply FARMALINX CLEARVIEW 200 GR Herbicide by uniformly distributing by hand after dispensing a single dose (7g) from the dispenser unit. Apply

www.farmalinx.com

CONTINUED OVERLEAF

RESISTANT WEEDS WARNING
 FARMALINX CLEARVIEW 200 GR HERBICIDE is a member of the Urea group of herbicides. CLEARVIEW 200 GR has inhibitors of photosynthesis at photosystem II mode of action. For weed resistance management CLEARVIEW 200 GR is a Group C herbicide. Some naturally occurring weed biotypes resistant to CLEARVIEW 200 GR and other Group C herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by CLEARVIEW 200 GR or other Group C herbicides.
 Since the occurrence of resistant weeds is difficult to detect prior to use, FARMALINX Pty Ltd accepts no liability for any losses that may result from the failure of CLEARVIEW 200 GR to control resistant weeds.
PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS
 DO NOT allow any roots of desirable plants, trees or ornamentals to contact FARMALINX CLEARVIEW 200 GR Herbicide. FARMALINX CLEARVIEW 200 GR Herbicide may cause temporary herbicidal symptoms to appear on perennial grasses, including buffer grass, particularly after extended dry periods. Higher rates of application of FARMALINX CLEARVIEW 200 GR Herbicide (2g/m²) may kill grasses, particularly native annuals. This effect

Treatment of Clumps of Trees or Patches of Regrowth
 FARMALINX CLEARVIEW 200 GR Herbicide pellets are particularly effective in treating areas of regrowth.
 Estimate the area within 30cm beyond the drip zone and apply the FARMALINX CLEARVIEW 200 GR Herbicide pellets on a grid pattern to uniformly cover the area. Pellets should be applied over the top of regrowth if at all possible in an attempt to maintain a good distribution pattern.
Example:


Treating Fence Lines
 Estimate the area to be treated, along the fence line and then apply FARMALINX CLEARVIEW 200 GR Herbicide pellets on both sides of the fence line as evenly as possible within the area extending 30cm out beyond drip zone.
Example:


the pellets at 1 to 2g per metre square. One dose (7g) will treat 7 to 3.5 square metres at these rates.
ADDITIONAL INSTRUCTIONS
 When applying pellets, distribute them under the tree or regrowth canopy from the root crown or stems to 30cm beyond the drip zone of the plants you want to control. The time required for complete plant death depends on soil type, amount of rainfall, root depth, plant species, plant density and rate of FARMALINX CLEARVIEW 200 GR Herbicide used. Some species may go through several defoliations from 6 months to 2 years before dying. Timber and regrowth control with FARMALINX CLEARVIEW 200 GR Herbicide pellets may be variable when used on soils with high clay content high organic matter or when treating deep rooted or more tolerant species.

TREATING INDIVIDUAL TREES
 Estimate the area within 30cm beyond the drip zone, i.e. the area under the plant canopy. It is essential that this area is treated as evenly as possible with pellets being spread from the tree trunk to slightly beyond the drip line formed by the branches.
 DO NOT throw all the pellets at the base of the tree trunk. This will waste material and could result in poor control.

Example:
