

FARMALINX

Flufol 600 SC

FUNGICIDE

DIRECTIONS FOR USE: ALL STATES

Restraints: DO NOT use open loading and mixing for aerial application.

A. FARMALINX FLUFOL 600 SC FUNGICIDE APPLIED AS A FERTILISER TREATMENT

Crop	Disease	Rate per hectare	WHP	Critical Comments				
Canola	Blackleg (<i>Leptosphaeria maculans</i>)	167 mL	4 weeks Grazing	Treatment will reduce blackleg infection in both susceptible and tolerant canola varieties. Application: Granulated Fertiliser Spray undiluted product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate, which will deposit 400mL of the product on the quantity of fertiliser required to be sown in furrow per hectare, using the table below as a guide.				
				Fertiliser rate per hectare (kg)	Volume of product per 100 kg of Granular Fertiliser (mL/100 kg)			
				60	278			
				70	238			
				80	208			
100	167 [†]							
120	139 [†]							
† Where application volume is low, product may be diluted with water to improve coverage.								
Wheat	Stripe rust (<i>Puccinia striiformis</i> f.sp. <i>tritici</i>)	83 to 333 mL	4 weeks Grazing	For stripe rust control in wheat use 83 to 167 mL/ha where up to 100 days protection is required and where heavy stripe rust pressure is expected use 167 mL/ha only. Use 250 to 333 mL/ha when 120 to 130 days protection is required and where heavy stripe rust pressure is expected, use 333 mL/ha only.				
	Septoria tritici blotch (<i>Mycosphaerella graminicola</i>) Take-all (<i>Geaumannomyces graminis</i>)	167 mL		For control of powdery mildew and scald of barley, use 83 to 167 mL/ha where up to 80 days protection is required and where heavy disease pressure is expected use 167 mL/ha. Application: Spray product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit the specific rate of the product on the quantity of fertiliser required to be sown in-furrow per hectare, using the table below as a guide.				
Barley	Powdery mildew (<i>Blumeria graminis</i> f.sp. <i>hordei</i>) Scald (<i>Rhynchosporium secalis</i>)	83 to 167 mL		Granular Fertiliser rate (kg/ha)	Volume of product per 100 kg of Granular Fertiliser (mL/100 kg)			
					83 mL rate	167 mL rate	250 mL rate (Wheat only)	333 mL rate (Wheat Only)
				50	167 [†]	333	500	667*
				60	139 [†]	278	417	556
				70	119 [†]	238	358	476
				80	104 [†]	208	313	417
				100	83 [†]	167 [†]	250	333
120	69 [†]	139 [†]	208	278				
* Application volumes over 600 mL per 100 kg of fertiliser may adversely affect the flow rate and handling characteristics of dusty/poorly granulated fertilisers and some high nitrate fertilisers. † Where application volume is low, product may be diluted with water to improve coverage.								

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

**A. APPLIED IN-FURROW AS A FERTILISER TREATMENT
CANOLA, WHEAT AND BARLEY - DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER SOWING**

GENERAL INSTRUCTIONS

Fertiliser Application

Granulated Fertiliser

To determine the correct volume of Flufol 600 SC required to be applied, refer to Critical Comments in the Directions for Use table, which provides a guide based on a range of common fertiliser application rates. Ensure that treated fertiliser is dry before sowing with the crop.

Treated fertiliser should be applied in a band 3-4 cm below or to the side of seed in planting furrow. It should not be broadcast. Note that the flow rate of fertiliser may be changed after treatment, and the seeder should be calibrated with the treated material to ensure that the correct rates of fertiliser and fungicide are delivered. Do not mix seed with treated fertilizer and Flufol 600 SC.

Re-entry or Re-handling

DO NOT re-handle treated fertiliser until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves.

FUNGICIDE RESISTANCE WARNING

Farmalinx Flufol 600 SC Fungicide is a member of the DMI group of fungicides. For fungicide resistance management Flufol 600 SC Fungicide is a Group C fungicide. Some naturally occurring individual fungi resistant to Flufol 600 SC Fungicide and other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Flufol 600 SC Fungicide or other Group C fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Farmalinx Pty Ltd accepts no liability for any losses that may result from the failure of Flufol 600 SC Fungicide to control resistant fungi. Resistance should not be assumed without first reviewing the method of application, the coverage and the timing of application.

GROUP 3 FUNGICIDE

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

SDS

Additional information is listed in the safety data sheet (SDS). A safety data sheet for Farmalinx Flufol 600 SC Fungicide is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 9389 2455.

NOTICE:

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Farmalinx Pty Ltd or under abnormal conditions. Farmalinx Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

farmalinx
Partners in Agriculture

FARMALINX Pty Ltd ABN 95 134 353 245
Level 25, Suite 2506, Tower 2,
101 Grafton Street, Bondi Junction NSW 2022
T: 02 9389 2455 F: 02 9389 2844
Website: www.farmalinx.com

APVMA Approval NO.: 84898/111715

STORAGE AND DISPOSAL (10 – 20L)

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

STORAGE AND DISPOSAL (100 -1000L)

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs Farmalinx Pty Ltd should be advised immediately.

This container is reusable and remains the property of Farmalinx Pty Ltd. DO NOT rinse empty container. Empty contents fully into application equipment. Close all valves and return to the point of supply for refill or storage. No other liquid, solid or pesticide product should be put into it. When empty return to Farmalinx Pty Ltd for cleaning, relabelling and refilling.

SAFETY DIRECTIONS

Will irritate the eyes. May irritate the skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When using the product wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow length chemical resistant gloves and face shield or goggles. If product in eyes wash it out immediately with water. Wash hands after use. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Flufol 600 SC

FUNGICIDE

ACTIVE CONSTITUENT: 600g/L FLUTRIAFOL

GROUP 3 FUNGICIDE

For control of certain fungal diseases on wheat, barley and canola when mixed with fertiliser as per the Directions for Use

IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT

farmalinx

FARMALINX Pty Ltd ABN 95 134 353 245
Level 25, Suite 2506, Tower 2, 101 Grafton Street,
Bondi Junction NSW 2022