

# Firefox 400 SL

HERBICIDE

## DIRECTIONS FOR USE

### RESTRAINTS:

DO NOT apply by aircraft (except for sugarcane).

DO NOT apply if rain is expected within 3 hours (24 hours for bracken).

CROP	WEEDS CONTROLLED	RATE	STATE	CRITICAL COMMENTS
Apple orchards	Docks	<b>Boom sprayer</b> 4.2 L in 220 to 440 L water/ha <b>Knapsack or Hand lance</b> 1 part to 200 parts water	NSW, Vic, Tas, SA, WA only	Apply when docks are actively growing and have fully expanded leaves, just before flowering shoots emerge. With knapsack or hand lance, ensure uniform coverage but do not spray to run-off. Apply only to established trees. DO NOT graze the area to be sprayed for 7 days before and 21 days after application.
Hops	Broadleaf dock, curled dock, fiddle dock	3.5 to 5.0 L in 220 to 440 L water/ha	Vic, Tas only	Apply before the hop plants have emerged. Apply the higher rate where fiddle dock is present or where docks are mainly old, established plants.
Lucerne - established	Docks	<b>Boom sprayer</b> 2.8 to 4.2 L in 220 to 440 L water/ha	All states	Lucerne should be at least one year old. Apply when docks are actively growing and have fully expanded leaves, just before flowering shoots emerge. Apply lower rate if curled dock and broadleaf dock are present. Apply higher rate if other docks are present. DO NOT apply in spring if pasture is intended for seed production.
Onions	Docks	<b>Knapsack or Hand lance</b> 1 part to 200 parts water	Tas only	Apply when docks are actively growing and have fully expanded leaves, just before flowering shoots emerge. Ensure uniform spray coverage of docks but do not spray to run-off. DO NOT apply as an overall spray as crop damage may result. Sprayed onions may show a slight yellowing but they will normally recover quickly if good growing conditions exist.
Oilseed poppies	Broadleaf dock, curled dock	2.5 L in 220 to 440 L water/ha	Tas only	<b>Crop stage:</b> spray any time after the 4 to 6 leaf stage. <b>Weed stage:</b> apply when docks are actively growing and have fully expanded leaves, just before flowering shoots emerge. The range of control is extended with a spray program in combination with Ethofumesate as recommended by poppy contracting companies or the Department of Primary Industries, Tasmania.
Pastures established & improved pastures of: Chewing's fescue, cocksfoot, perennial ryegrass, red clover, tall fescue, white clover	Common bracken	<b>Boom sprayer</b> *12.0 L/ha plus Bioclear Paraffinic Oil (2% of spray volume) in 110 to 330 L water/ha <b>Motorised mister</b> *12.0 L plus Bioclear Paraffinic Oil (2% of spray volume) in 20 to 80 L water/ha <b>Knapsack or hand lance</b> *100 mL plus Bioclear Paraffinic Oil (2% of spray volume) in 9.8 L water	All states	*Note the addition of a spray oil, may improve performance, however, temporary scorch is likely to occur in perennial pastures. Consult your local agronomist for advice.  Apply during late April to early June after the Autumn re-growth has fully expanded, and before the first frosts of the season. Where old mature fronds exist, the bracken should be slashed or burned during October or November in the year prior to spraying. Results are first seen as a lack of spring re-growth. Ensure uniform coverage of spray over bracken. With knapsack or hand lance, spray to give complete coverage of fronds but do not spray to run-off. DO NOT spray if rain is likely within 24 hours.
	Common bracken	<b>Carpet wiper</b> *5.0 L plus Bioclear Paraffinic Oil (2% of spray volume) /ha	Vic only	*Note the addition of a spray oil may improve performance, however, temporary scorch is likely to occur in perennial pastures. Consult your local agronomist for advice. For East Gippsland district only. To apply the recommended rate it is necessary to select an appropriate combination of flow rate, tractor speed and herbicide dilution. Refer to General Instructions for details. This rate will control bracken growing at a density of 25 fronds per square metre. The bracken should be slashed during late spring or early summer in the year prior to spraying to ensure regenerated fronds of similar age. The bracken should then be wiped in the following autumn before the first frosts of the season when the regenerated fronds are fully mature. Where fronds are of variable height or occur in dense infestation of clumps, some plants may not be contacted by the herbicide solution. In these cases, repeat treatment may be necessary.

CROP	WEEDS CONTROLLED	RATE	STATE	CRITICAL COMMENTS
Pastures established & improved pastures of: Chewing's fescue, cocksfoot, perennial ryegrass, red clover, tall fescue, white clover	Broadleaf dock, curled dock	<b>Boom sprayer</b> 2.8 L in 220 to 440 L water/ha	All states	Apply when docks are actively growing and have fully expanded leaves, just before flowering shoots emerge. Slight yellowing of some pasture species may occur but they will normally recover quickly if good growing conditions exist. Damage to strawberry clover and subterranean clover is likely. A follow-up spray may be required to control subsequent germinations of seedlings which occur in old established stands of mature docks. Pastures should not be grazed for 2 to 3 weeks before spraying to allow sufficient growth of docks and to minimize the check to pastures. DO NOT apply in spring if pasture is intended for seed production.
Perennial ryegrass for seed production	<b>Group A</b> Dock*, false brome, rat's tail fescue, shivery grass, silvery hair grass, soft brome, squirrel tail fescue, sub clover <b>Group B</b> Madrid brome, prairie grass, sorrel, sterile brome, annual poa (winter grass), Yorkshire fog	5.5 L in 160 to 220 L water/ha	NSW, Vic, Tas, SA only	Apply to established crops in the Autumn (late April-early June). The crop will show signs of yellowing 5-6 weeks after spraying. A reduction in total herbage production for 3-6 months will be experienced. Annual clovers will be affected. DO NOT apply to cultivar Medea ryegrass as crop yields may be affected. DO NOT apply in spring as crop damage will result. Group A weeds are susceptible and Group B weeds are moderately susceptible at the recommended rate. *Apply when docks are actively growing and have fully expanded leaves, just before flowering shoots emerge.
Potatoes	Docks	<b>Boom sprayer</b> 2.8 L in 220 to 440 L water/ha	Vic, Tas only	Apply when docks are actively growing and have fully expanded leaves, just before flowering shoots emerge. Apply before the potato foliage obscures the docks.
Sugar cane	Awnless barnyard grass, barnyard grass, crab grass, crowsfoot grass, green summer grass, giant paspalum, hairy panic itch grass, Johnson grass, para grass, summer grass	<b>Boom sprayer</b> 8.5 L in 200 to 400 L water/ha <b>Aircraft</b> 8.5 L in a min. 22 L water/ha <b>Spot application*</b> 200 mL in 9.8 L of water. In mixtures with Actril* DS use at least 55 L water/ha	Qld, NSW only	<b>Crop stage:</b> apply post-emergence in plant or ratoon cane. <b>Weed stage:</b> apply when weeds are actively growing, before flowering and before they exceed 200 to 250 mm in height unless otherwise indicated. Optimal results will be obtained when the relative humidity is high (over 80%) and temperature above 17°C. Residual control of grasses germinating from seed will persist for 6 to 10 weeks, depending on field conditions. <b>Surfactant:</b> Wetdrop Wetter at 200 mL/100 L to improve wettability and coverage and reduce spray runoff. <b>Directed application:</b> ideally the spray should be directed to minimize coverage of the cane foliage. Aircraft application may be necessary in salvage situations and could result in some crop damage. DO NOT apply tank mixes from aircraft. <b>Caution:</b> certain varieties have not been screened for tolerance to this product. Treat a small area first, by the same technique as that to be used commercially to check tolerance. Variety Q68 should not be treated. *With knapsack or hand lance, ensure uniform coverage but do not spray to run-off.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

### WITHOLDING PERIODS

**APPLE ORCHARDS, LUCERNE, ONIONS, PASTURES, PERENNIAL RYEGRASS:  
DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 21 DAYS AFTER APPLICATION**

www.farmalinx.com

CONTINUED OVERLEAF

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**  
DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

**STORAGE AND DISPOSAL**  
KEEP OUT OF REACH OF CHILDREN. Store in the closed original container in a dry, well-ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight. Wash the spraying equipment thoroughly before and after use. Triple or (preferably) pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycle or designated collection point. If not recycling, break, crush or puncture empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulation. Do not burn empty containers or product.

**SAFETY DIRECTIONS**  
Avoid contact with eyes and skin. DO NOT inhale spray mist. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

**FIRST AID**  
If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126.

**MATERIAL SAFETY DATA SHEET**  
Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FARMALINX FIREFOX 400 SL Herbicide is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 9389 2455.

Other trademarks

**NOTICE TO BUYER**  
Seller warrants that the product conforms to its chemical description and its reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off-label permits not endorsed by FARMALINX Pty Ltd or under abnormal conditions.

from incorrect storage, handling or use.

**farmalinx**  
Partners in Agriculture

FARMALINX Pty Ltd ABN 95 134 353 245  
Level 25, Suite 2506, Tower 2,  
101 Grafton Street, Bondi Junction NSW 2022  
T: 02 9389 2455 F: 02 9389 2844  
Website: www.farmalinx.com

APVMA Approval No. 69008/59635

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FARMALINX  
**Firefox 400 SL**

HERBICIDE

ACTIVE CONSTITUENT: 400 g/L ASULAM present as the SODIUM SALT

GROUP R HERBICIDE

*For control of bracken fern in pastures: grasses in sugar cane, and a range of weeds and grasses in crops and pastures as specified in the Directions for Use table.*

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.

FARMALINX Pty Ltd ABN 95 134 353 245  
Level 25, Suite 2506, Tower 2, 101 Grafton Street,  
Bondi Junction NSW 2022

**farmalinx**