

DIRECTIONS FOR USE:

A. FARMALINX FLUFOL APPLIED IN-FURROW AS A FERTILISER TREATMENT

CROP	DISEASE	RATE PER HECTARE	WHP	CRITICAL COMMENTS																					
Canola	Blackleg (<i>Leptosphaeria maculans</i>)	400 mL	4 weeks grazing	Treatment will reduce blackleg infection in both susceptible and tolerant canola varieties. Application: Spray undiluted product evenly onto granulated fertilizer and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate, which will deposit 400 mL of the product on the quantity of fertilizer required to be sown in furrow per hectare, using the table below as a guide. <table border="1"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th>Volume of product per 100 kg of fertiliser (mL)</th> </tr> </thead> <tbody> <tr><td>60</td><td>667*</td></tr> <tr><td>70</td><td>571</td></tr> <tr><td>80</td><td>500</td></tr> <tr><td>100</td><td>400</td></tr> <tr><td>120</td><td>333</td></tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)	60	667*	70	571	80	500	100	400	120	333									
Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)																								
60	667*																								
70	571																								
80	500																								
100	400																								
120	333																								
Wheat	Stripe Rust (<i>Puccinia striiformis</i>)	200 or 400 mL		For control of stripe rust of wheat and control of powdery mildew and scald of barley, use the high rate where conditions conducive to severe disease are expected. Application: Spray product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit the specific rate of the product on the quantity of fertiliser required to be sown in-furrow per hectare, using the table below as a guide. <table border="1"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th>Volume of product per 100 kg of fertiliser (mL)</th> </tr> </thead> <tbody> <tr><td>50</td><td>400</td></tr> <tr><td>60</td><td>333</td></tr> <tr><td>70</td><td>286</td></tr> <tr><td>80</td><td>250#</td></tr> <tr><td>100</td><td>200#</td></tr> <tr><td>120</td><td>167#</td></tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)	50	400	60	333	70	286	80	250#	100	200#	120	167#							
	Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)																							
50	400																								
60	333																								
70	286																								
80	250#																								
100	200#																								
120	167#																								
	Septoria Tritici Blotch (<i>S. tritici</i>), Take-all (<i>Geaumannomyces graminis</i>)	400 mL																							
Barley	Powdery Mildew (<i>Erysiphe graminis</i>), Scald (<i>Rhynchosporium secalis</i>)	200 or 400 mL		<table border="1"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th>200 mL rate</th> <th>400 mL rate</th> </tr> </thead> <tbody> <tr><td>50</td><td>400</td><td>800*</td></tr> <tr><td>60</td><td>333</td><td>667*</td></tr> <tr><td>70</td><td>286</td><td>571</td></tr> <tr><td>80</td><td>250#</td><td>500</td></tr> <tr><td>100</td><td>200#</td><td>400</td></tr> <tr><td>120</td><td>167#</td><td>333</td></tr> </tbody> </table> * Application volumes over 600 mL per 100 kg of fertiliser may adversely affect the flow rate and handling characteristics of dusty/poorly granulated fertilisers and some high nitrate fertilisers. # Where application volume is low, product may be diluted with water to improve coverage.	Fertiliser rate per hectare (kg)	200 mL rate	400 mL rate	50	400	800*	60	333	667*	70	286	571	80	250#	500	100	200#	400	120	167#	333
Fertiliser rate per hectare (kg)	200 mL rate	400 mL rate																							
50	400	800*																							
60	333	667*																							
70	286	571																							
80	250#	500																							
100	200#	400																							
120	167#	333																							

B. FARMALINX FLUFOL APPLIED AS A FOLIAR SPRAY

CROP	DISEASE	RATE PER HECTARE	WHP	CRITICAL COMMENTS
Wheat	Stripe Rust (<i>Puccinia striiformis</i>)	250-500 mL Ground Application Apply in 100-150 L water plus 200 mL FARMALINX WetDrop per 100 L of spray solution.	7 weeks Harvest or Grazing	Apply when stripe rust appears and before disease reaches 1% of leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with high yield potential and where conditions favourable to severe disease persist, use a single application at the higher rate, or two applications 3-4 weeks apart at the low rate.
	Leaf Rust (<i>P. recondita</i>), Septoria Nodorum Blotch (<i>Septoria nodorum</i>), Septoria Tritici Blotch (<i>S. tritici</i>)	Aerial Application Apply in a minimum of 15 L water plus 200 mL FARMALINX WetDrop per 100 L of spray solution.		Apply after flag leaves have begun to emerge and before disease reaches 1% of flag leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with high yield potential and where conditions favourable to severe disease persist during flowering, use a single application at the higher rate, or two applications 3-4 weeks apart at the low rate.
Barley	Powdery Mildew (<i>Erysiphe graminis</i>)	100 L of spray solution.	10 weeks Harvest or Grazing	Apply when powdery mildew first appears. The high rate of application gives a longer period of protection than the low rate. Where conditions favourable to severe disease persist, a single application at the higher rate, or two applications 3-4 weeks apart at the low rate, may be required.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

APPLIED IN-FURROW AS A FERTILISER TREATMENT

CANOLA, WHEAT AND BARLEY: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER SOWING.

APPLIED AS A FOLIAR SPRAY

DO NOT HARVEST EARLIER THAN THE NUMBER OF WEEKS SHOWN AFTER APPLICATION:

BARLEY – 10 WEEKS

WHEAT – 7 WEEKS

DO NOT GRAZE OR CUT FOR STOCK FOOD EARLIER THAN THE NUMBER OF WEEKS SHOWN AFTER APPLICATION:

BARLEY – 10 WEEKS

WHEAT – 7 WEEKS

GENERAL INSTRUCTIONS

FUNGICIDE RESISTANCE WARNING

FARMALINX Flufol In-Furrow & Foliar Fungicide is a member of the DMI group of fungicides. For fungicide resistance management FARMALINX Flufol In-Furrow & Foliar Fungicide is a Group 3 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled this product and other Group 3 fungicides thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, FARMALINX Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi. Resistance should not be assumed without first reviewing the method of application, the coverage and the timing of application.

APPLICATION

Fertiliser Application

Ensure that treated fertiliser is dry before sowing with the crop.

Sowing: Treated fertiliser should be sown in the same furrow as the seed, not broadcast. Note that the flow rate of fertiliser may be changed after treatment, and the seeder should be calibrated with the treated material to ensure that the correct rates of fertiliser and fungicide are delivered. DO NOT mix seed with treated fertiliser.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre.

Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FARMALINX Flufol In Furrow & Foliar Fungicide is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 9389 2455.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off-label permits not endorsed by FARMALINX Pty Ltd or under abnormal conditions. FARMALINX Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

farmalinx
Partners in Agriculture

FARMALINX Pty Ltd ABN 95 134 353 245

Level 25, Suite 2506, Tower 2,

101 Grafton Street, Bondi Junction NSW 2022

T: 02 9389 2455 F: 02 9389 2844

Website: www.farmalinx.com

APVMA Approval No.: 64291/0809

GROUP 3 FUNGICIDE

Foliar Application

Check the drop carefully and regularly. Spray only if there is sufficient time in the season for the rust to develop to a level on the variety in question when it will cause significant yield loss.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

The method of disposal of the container depends on the container type.

Read the storage and disposal instructions on the label that is attached to the container.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. May irritate the nose and throat and skin. Avoid contact with eyes and skin. DO NOT inhale vapour. When opening the container and preparing the spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and face shield or goggles. If product in eyes, wash out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FARMALINX

Flufol

IN FURROW & FOLIAR FUNGICIDE

ACTIVE CONSTITUENT: 250 g/L FLUTRIAFOL

GROUP 3 FUNGICIDE

For the control of certain fungal diseases on wheat, barley and canola when mixed with fertiliser or applied as a foliar spray as per the Directions for Use Table.

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT

FARMALINX Pty Ltd ABN 95 134 353 245
Level 25, Suite 2506, Tower 2, 101 Grafton Street,
Bondi Junction NSW 2022

farmalinx