

FARMALINX

Holt 750 WG

HERBICIDE

DIRECTIONS FOR USE:

RESTRAINTS:

DO NOT APPLY THIS PRODUCT WHEN TEMPERATURES ARE GREATER THAN 18°C FOR THE DAY PRIOR TO, THE DAY OF AND FOR 4 DAYS FOLLOWING APPLICATION.

DO NOT spray under dry frosty conditions or if very dry conditions prevail.

DO NOT spray if weeds or crop/pastures are affected by insects, disease, dust or adverse weather conditions.

DO NOT add wetting agents or crop oils.

DO NOT use main or nozzle filters finer than 50 mesh for tank mixes.

DO NOT tank mix with selective grass herbicides.

DO NOT apply tank mixture of FARMALINX HOLT 750 WG Herbicide plus triasulfuron (750 g/kg) WG if another Group B herbicide (ASL inhibitor) has been used during the current season.

IF RAIN FALLS within 6 hours of application, the effect could be diminished.

CROP	WEEDS CONTROLLED	STATE	RATE/ha	CRITICAL COMMENTS
Wheat, Triticale, Barley, some varieties of Oats: from 3 leaf to early tillering stage	Amsinckia (Yellow Burrweed), Capeweed, Corn Gromwell (Sheep Weed or White Iron Weed), Deadnettle, Fumitory (Common, White and Dense Flowered), Paterson's Curse, Toad Rush – 2 to 6 leaves	NSW, Vic only	367 to 567 g	Use the highest rate when Deadnettle or Toad Rush is the major problem. Apply to boom spray or aircraft.
	The above weeds in addition to Charlock, Doublegee (Three-cornered Jack), Hedge Mustard, Hexham Scent, Indian Hedge Mustard, Turnip Weed, Wild Turnip – 2 to 6 leaves		367 to 567 g + 300 mL of a 500 g/L 2,4-D (as amine) or 300 mL of a 500 g/L MCPA as a tank mix	DO NOT add MCPA or 2,4-D if undersown Medics are present. Apply by boom spray or aircraft.
	<i>Plagiobothrys canascens</i> – 2 to 6 leaves		567 g	Use the highest rate when Deadnettle, Doublegee (Three-cornered Jack) or Toad Rush is the major problem.
	Turnip Weed	NSW only	200 g + 6.5 g of TRIAFON	Spray when weeds are in the 2 to 6 leaf stage (up to 6 node/leaflet for Field Peas, 10 leaflet for Vetch), except 2 to 4 leaf for Doublegee.
	Hedge Mustard, Indian Hedge Mustard, London Rocket, Wild Turnip		200 g + 6.5 g of TRIAFON to 293 g+ 10 g of TRIAFON	
	Black Bindweed (Climbing Buckwheat), Faba Bean (volunteer), Field Pea (volunteer), Medic, Shepherd's Purse		293 g+ 10 g of TRIAFON	Use rates towards the lower end of the range (where applicable) when weeds are small and soil conditions are very moist.
	Deadnettle, Mexican Poppy, Wild Radish		293 g+ 10 g of TRIAFON to 400 g + 13 g of TRIAFON	Spray only when weeds are actively growing.
	Coreopsis, Sunflower (volunteer), Vetch		400 g + 13 g of TRIAFON	Spray only after good rain and when top soil is moist.
	Paterson's Curse, Turnip Weed	Vic only	200 g + 6.5 g of TRIAFON	Best results are obtained when good soil moisture has been present since planting.
	Hedge Mustard, Indian Hedge Mustard, Wild Turnip		200 g + 6.5 g of TRIAFON to 293 g+ 10 g of TRIAFON	
Crassula, Fabia Bean (volunteer), Hyssop Loosetrife, Medic, Prickly Lettuce (Whipthistle)	293 g+ 10 g of TRIAFON			
Deadnettle, Field Pea (volunteer), Wild Radish	293 g+ 10 g of TRIAFON to 400 g + 13 g of TRIAFON			
Lupins (volunteer), Vetch, Wireweed (less than 3 leaves) suppression	400 g + 13 g of TRIAFON			

CROP	WEEDS CONTROLLED	STATE	RATE/ha	CRITICAL COMMENTS	
Wheat, Triticale, Barley, some varieties of Oats: from 3 leaf to early tillering stage <i>cont.</i>	Turnip Weed	Qld only	200 g + 6.5 g of TRIAFON	Spray when weeds are in the 2 to 6 leaf stage, except 2 to 4 leaf for Doublegee.	
	Denseflower Fumitory, Indian Hedge Mustard, Smallflower Fumitory, Wild Turnip		200 g + 6.5 g of TRIAFON to 293 g+ 10 g of TRIAFON	Use rates towards the lower end of the range (where applicable) when weeds are small and soil conditions are very moist.	
	Black Bindweed (Climbing Buckwheat), London Rocket		293 g+ 10 g of TRIAFON	Spray only when weeds are actively growing.	
	Wild Radish		293 g+ 10 g of TRIAFON to 400 g + 13 g of TRIAFON	Best results are obtained when good soil moisture has been present since planting.	
	Coreopsis, Corn Gromwell (Sheepweed or White Ironweed)		400 g + 13 g of TRIAFON		
		Amsinckia (Yellow Burrweed), Capeweed, Corn Gromwell (Sheep Weed or White Iron Weed), Deadnettle, Doublegee (Three-cornered Jack), Fumitory (Common, White and Dense Flowered), Rough Poppy, Salvation Jane, Toad Rush – 2 to 6 leaves	SA only	367 to 567 g	DO NOT add MCPA or 2,4-D if undersown Medics are present. Apply by boom spray or aircraft. Use the highest rate when Deadnettle, Doublegee (Three-cornered Jack) or Toad Rush is the major problem.
		The above weeds in addition to Charlock, Hedge Mustard, Hexham Scent, Indian Hedge Mustard, Mallee Catch-Fly, Turnip Weed, Wards Weed, Wild Turnip – 2 to 6 leaves		367 to 567 g + 300 mL of a 500 g/L 2,4-D (as amine) or 300 mL of a 500 g/L MCPA as a tank mix	DO NOT apply to disease infected wheat as yellowing may occur.
		Faba Beans (volunteer), Long Fruited Turnip, Long Headed Poppy, Paterson's Curse, Turnip Weed		200 g + 6.5 g of TRIAFON	Spray when weeds are in the 2 to 6 leaf stage (up to 6 node/leaflet for Field Peas, 10 leaflet for Vetch), except 2 to 4 leaf for Doublegee. Use rates towards the lower end of the range (where applicable) when weeds are small and soil conditions are very moist. Spray only when weeds are actively growing. Spray only after good rain and when top soil is moist. Best results are obtained when good soil moisture has been present since planting.
		Hedge Mustard, Indian Hedge Mustard, Wild Turnip		200 g + 6.5 g of TRIAFON to 293 g+ 10 g of TRIAFON	
		Ball Mustard, Crassula, Medic, Prickly Lettuce (Whipthistle)		293 g+ 10 g of TRIAFON	
Field Pea (volunteer), Wild Radish		293 g+ 10 g of TRIAFON to 400 g + 13 g of TRIAFON			
Doublegee (Spiny Emex, Three-cornered Jack), Lupins (volunteer), Vetch, Wireweed (less than 3 leaves) suppression		400 g + 13 g of TRIAFON			
Capeweed, Corn Gromwell, Fumitory, Yellow Burrweed – 2 to 6 true leaves, Doublegee (Three-cornered Jack), Toad Rush – 1 to 4 leaves		WA only		567 g	
The above weeds in addition to Indian Hedge Mustard, London Rocket, Lupin, Wild Radish and Wild Turnip – 2 to 6 leaves				367 to 567 g or 600 mL of a 500 g/L MCPA as a tank mix	Use the highest rate when Fumitory is the major problem weed in addition to Wild Turnip, Indian Hedge Mustard, London Rocket, Lupin and Wild Radish.
Turnip Weed				200 g + 6.5 g of TRIAFON	Spray when weeds are in the 2 to 6 leaf stage (up to 6 node/leaflet for Field Peas, 10 leaflet for Vetch), except 2 to 4 leaf for Doublegee. Use rates towards the lower end of the range (where applicable) when weeds are small and soil conditions are very moist. Spray only when weeds are actively growing. Spray only after good rain and when top soil is moist.
Denseflower Fumitory, Hedge Mustard, Smallflower Fumitory	200 g + 6.5 g of TRIAFON to 293 g+ 10 g of TRIAFON				
Ball Mustard, Field Pea (volunteer)		293 g+ 10 g of TRIAFON			
Deadnettle		293 g+ 10 g of TRIAFON to 400 g + 13 g of TRIAFON			
Australian Crassula, Doublegee (Spiny Emex, Three-cornered Jack), Lupins (volunteer)		400 g + 13 g of TRIAFON	Best results are obtained when good soil moisture has been present since planting.		

CROP	WEEDS CONTROLLED	STATE	RATE/ha	CRITICAL COMMENTS
Pastures Use on mixtures of either/or Sub Clover (min. of 3 to max. of 10 leaves), Balansa Clover (established plants), White Clover (min. of 5 to max. of 10 leaves) with either/ or Ryegrass, Phalaris, Cocksfoot (min. 3 leaves)	Capeweed (seedling to 10 cm diameter), Paterson's Curse (seedling to 10 cm diameter)	NSW, Vic, Tas only	367 to 467 g	Apply by boom spray or aircraft. Use the lowest rate early in the season or when weeds are small and young. Increase rate as weed size and age increases.
	Capeweed (seedling to 20 cm diameter), Crassula (1 to 5 cm diameter), <i>Erodium botrys</i> (seedlings to 8 cm diameter), Paterson's Curse (seedling to 20 cm diameter), Toad Rush (1 to 5 leaves)		267 g to 467 g plus 300 to 500 mL of MCPA 500 amine as a tank mix	Early germinating Paterson's Curse with strong root systems may not be controlled. Some temporary damage may be seen on clover if adverse growing conditions occur at end after spraying. Where a selective grass herbicide or simazine is to be or has been used. FARMALINX HOLT 750 WG Herbicide should not be used in the same season. See Pasture Safety section in General Instructions.
Field Peas	Amsinckia (Yellow Burrweed), Capeweed, Common Fumitory, Corn Gromwell (Sheep Weed or White Iron Weed), Deadnettle, Paterson's Curse, Toad Rush	NSW, Vic, Tas, SA only	467 to 567 g	Apply as an early post-emergent application when the peas are 10 to 12 cm high and have developed 3 to 5 nodes. Weeds should be young and actively growing and in the 4 to 5 true leaf stage. Use rates towards the higher end of the scale when Deadnettle is the major problem and when weeds are in the 6 true leaf stage. Where a selective grass herbicide is to be used, apply FARMALINX HOLT 750 WG Herbicide 2 weeks before the application of the selective herbicide.
Canning Peas (Green Peas, Vining Peas, Processing Peas)	Blackberry Nightshade, Capeweed, Chickweed, Common Fumitory, Corn Gromwell, Deadnettle, Fat Hen, Paterson's Curse, Toad Rush, Yellow Burrweed	Tas only	500 to 667 g	Apply as an early post-emergent application when the peas are 10 to 12 cm high and have developed 3 to 6 nodes and the weeds are young and actively growing and in the 2 to 4 true leaf stage. Use rates towards the higher end of the scale when Deadnettle and Blackberry Nightshade are the major problems.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THE LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHOLDING PERIODS:

FIELD PEAS, CANNING PEAS: DO NOT APPLY WITHIN 4 WEEKS OF HARVEST.

PEA FODDER AND FORAGE: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

PASTURES: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

CEREALS: FARMALINX HOLT 750 WG HERBICIDE ALONE AND FARMALINX HOLT 750 WG HERBICIDE PLUS MCPA OR

2,4-D AS A TANK MIX: DO NOT CUT OR GRAZE FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

CEREALS: FARMALINX HOLT 750 WG HERBICIDE PLUS TRIASULFURON WG TANK MIX: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Important

- Spray only after good rain and when topsoil is moist and when weeds are actively growing.
- Best results are obtained when good soil moisture has been present since planting.
- DO NOT apply during dry or waterlogged conditions.

RESISTANT WEEDS WARNING

FARMALINX HOLT 750 WG Herbicide is a member of the



triazine group of herbicides and has the photosystem II photosynthesis inhibitor mode of action. For weed resistance management FARMALINX HOLT 750 WG Herbicide is a Group C herbicide. Some naturally occurring weed biotypes resistant to FARMALINX HOLT 750 WG Herbicide and other Group C herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by FARMALINX HOLT 750 WG Herbicide or other Group C herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, FARMALINX Pty Ltd accepts no liability for any losses that may result from the failure of FARMALINX HOLT 750 WG Herbicide to control resistant weeds.

Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or a FARMALINX representative.

CROP SAFETY

DO NOT apply this product to Avon, Saia, Barmah and Cassia varieties of oats. Before applying to other varieties of oats, growers should seek advice from FARMALINX Pty Ltd for crop safety information. Undersown Medics and American Lucerne varieties may be damaged if sprayed. With early and late sown crops damage may occur if spraying is undertaken when temperatures are greater than 18°C. Avoid spraying crops prior to May and from mid-August onwards as daily temperatures and weed size may not be satisfactory. Early sown crops should not be sprayed before the onset of frost and cooler weather.

PASTURE SAFETY

DO NOT apply this product to pasture species other than those stated in the Directions for Use.

DO NOT apply to medics and lucerne as damage may occur.

DO NOT apply to irrigated pasture or seed crop as damage may occur.

DO NOT apply on pastures prior to May or after mid-August or when temperatures are greater than 18°C for the day prior to, the day of and for 4 days following application as pasture damage may occur.

DO NOT apply prior to or following the use of a grass selective herbicide or simazine in the same season as damage to pasture may occur.

DO NOT use in situations where oversowing or pastures may occur.

Pastures which contain young species should be sprayed using the lower label rates. Increase product rate as pasture age and size increases.

Only apply this product when conditions at and following application are favourable to pasture growth.

MIXING

Partly fill the spray tank with water and add the required amount of product while stirring. If required, add compatible products and agitate thoroughly. Continue agitation as filling of the tank is completed.

Agitate again before spraying commences.

If tank mixing with other water dispersible granule (WG) or wettable powder (WP) formulations, ensure they are mixed in well before adding emulsifiable concentrate (EC) or suspension concentration (SC) products. WP formulations should be pre-mixed separately prior to adding to the spray tank.

APPLICATION

Ground Application: Boom spray with flat fan nozzles. Ensure good spray coverage is obtained. Aim to apply 50 to 100 L per hectare.

Aircraft Application: Always spray in a cross wind of less than 5 knots, or use smoke generator to indicate air movements. Ensure good spray coverage is obtained. For maximum coverage use fine nozzles. Apply 20 to 30 L per hectare.

CROP ROTATION GUIDELINES

FARMALINX HOLT 750 WG Herbicide plus TRIAFON (triasulfuron (750 g/kg) WG) tank mixture

The following crops can be planted after:

In areas where pH is less than 7.5

– 9 months after an application: barley, cereal rye, chickpeas, faba beans, field peas, lupins, medics, oats, canola, safflower, subclover, triticale, wheat.

In areas where pH is greater than 7.5

– 9 months after an application: barley, cereal rye, oats, triticale, wheat.

– 14 months after an application: cotton, maize, sorghum, soybeans, sunflowers. Where residual herbicides are applied following an application of FARMALINX HOLT 750 WG Herbicide plus triasulfuron (750 g/kg) WG, the crop rotational guidelines for these products must be followed.

– 22 months after an application: chickpeas, faba beans, field peas, lupins, medics, canola, safflower, subclover.

For all other crops advice should be sought from FARMALINX Pty Ltd.

COMPATIBILITY

This product is compatible with 2,4-D Amine, MCPA 500 Amine (not LVE), triasulfuron (750 g/kg) WG (FARMALINX Triafon), FARMALINX Dicam, Dicamba + MCPA, Metsulfuron-Methyl (600 g/kg) WG (FARMALINX Metfuron) and Methidathion (400 g/L) EC.

PRECAUTION

Re-entry period: DO NOT enter treated area until spray has dried.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT use in situations where off-target damage may occur, eg vines, tree crops. If using 2,4-D amine or MCPA apply with equipment which will not be used in hormone susceptible crops, particularly cotton and vines. DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift over nearby susceptible plants or crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

1-25 kg HDPE Containers

Store in the closed, original container in a cool, well-ventilated area as cool as possible, out of direct sunlight. This container can be recycled if it is clean, dry, free of visible residues and has the *drumMUSTER* logo visible. Triple rinse container for disposal. Add rinsing to spray tank. DO NOT dispose of undiluted chemicals on-site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any *drumMUSTER* collection or similar container management program site. The cap should not be replaced but may be taken separately. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulation. DO NOT burn empty containers or product.

For refillable containers, empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

15 kg HDPE Bags

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight.

Single-rinse or shake remainder into spray tank. Do not dispose of undiluted chemicals on site. Puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Harmful if swallowed. May irritate the skin. Will irritate the eyes. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When preparing the product for use, wear elbow-length rubber gloves. Wash hands after use. After each day's use, wash gloves.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET

Additional information is listed in the safety data sheet (SDS). A material safety data sheet for FARMALINX HOLT 750 WG Herbicide is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 9389 2455.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Farmalinx Pty Ltd or under abnormal conditions. Farmalinx Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FARMALINX
Holt 750 WG

HERBICIDE

ACTIVE CONSTITUENT: 750 g/kg TERBUTRYN

GROUP C HERBICIDE

Controls certain early competing broadleaf weeds in wheat, barley, triticale, pastures, some varieties of oats and field peas as per Directions for Use table.

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT

farmalinx

FARMALINX Pty Ltd ABN 95 134 353 245
Level 25, Suite 2506, Tower 2, 101 Grafton Street,
Bondi Junction NSW 2022

farmalinx
Partners in Agriculture

FARMALINX Pty Ltd ABN 95 134 353 245

Level 25, Suite 2506, Tower 2

101 Grafton Street, Bondi Junction NSW 2022

T: 02 9389 2455 F: 02 9389 2844

Website: www.farmalinx.com

APVMA Approval No.: 83323/107978