

FARMALINX

# Metapris 500 SC

FUNGICIDE

**DIRECTIONS FOR USE:**  
**TREE AND VINE CROPS**

RATE					CRITICAL COMMENTS
In the following table, all rates are given for dilute spraying. For concentrate spraying, refer to the Mixing/Application section					
CROP	DISEASE	STATE	RATE	WHP	
Grapevines (wine grapes only)	Grey Mould ( <i>Botrytis cinerea</i> )	All States	75 mL/100 L	9 days	DO NOT use on table grapes or grapes used for the production of dried fruit. Use on wine grapes only. Apply at the following growth stages: 1. 80% cap fall. 2. Just prior to bunch closure. 3. At veraison (when sugar content rises). 4. 2-3 weeks pre-harvest. To ensure complete bunch wetting add "Agral" at 10-20 mL/100 L. DO NOT apply at concentrate rates greater than 150 mL/100 L.
Stone Fruit	Blossom Bight ( <i>Monilinia laxa</i> )	Qld, NSW, Vic, Tas, SA only	50-75 mL /100 L		DO NOT APPLY AFTER SHUCK FALL. Apply at 10% blossom, full bloom, late petal and shuck fall. If weather conditions particularly favour Blossom Bight use higher rate. <b>NSW, SA, Qld and Vic only:</b> Where <i>Monilinia laxa</i> is known to occur apply an additional early spray at pink bud. DO NOT apply at concentrate rates greater than 150 mL/100 L.
Navy Beans	Sclerotinia Rot ( <i>Sclerotinia sclerotiorum</i> )	Qld, NSW, Vic, SA, WA, NT only	100-150 mL/ 100 L		Use 1000 L water/ha or 1-1.5 L/ha. Apply in sufficient water to provide good coverage. Spray when 75% of plants first show open blossom and again 7 days later. Use the higher rate when disease pressure is high.
Faba beans	Chocolate Spot ( <i>Botrytis fabae</i> )	NSW, Vic, Tas, SA, WA only	50 mL/ha Ground Application in 100 L/ha		Inspect crop regularly on the mid to lower leaves, where disease will occur first. Apply spray in the early stages of attack, i.e. 1-2 spots per leaflet and when weather conditions favour the disease. These are temperatures between 15 and 20°C (Tasmania 10-20°C) and moist humid conditions (66% RH and over). If these conditions recur, repeat applications at intervals of 2-3 weeks will be necessary to protect new growth.
Onions	White Rot ( <i>Sclerotinia cepivorum</i> )	Qld, NSW, Vic, SA, NT only	20 mL/kg seed	4 weeks	<b>Seed Treatment</b> a) Apply 20 mL of FARMALINX Metapris 500 SC Fungicide to 1 kg of seed and mix thoroughly until seeds are wet. b) Spread the seed and allow to dry. c) Sow within 14 days of treatment. <b>Note:</b> 1. Seed treatment should be used in conjunction with soil applications of FARMALINX Metapris 500 SC Fungicide to achieve satisfactory control of White rot of onions. 2. <b>Caution:</b> Treated seed germinated poorly in cold, wet soil. Where these conditions occur, use a soil spray without seed treatment. 3. <b>Warning:</b> Soil persistence of FARMALINX Metapris 500 SC Fungicide can be reduced under alkaline soil conditions.

CROP	DISEASE	STATE	RATE	WHP	CRITICAL COMMENTS
Onions - continued	White Rot ( <i>Sclerotinia cepivorum</i> )	Qld, NSW, Vic, Tas, SA only	4 L/ha	4 weeks	<b>In-Furrow Application:</b> a) Thoroughly mix 4 L of FARMALINX Metapris 500 SC Fungicide with required quantity of fertiliser for 1 hectare. b) Apply fertiliser in a band no more than 2 cm directly below seed. <b>Note:</b> 1. Coarse sand or fine gravel can be substituted where fertiliser is not required. 2. In-furrow application must be combined with seed treatment to achieve satisfactory results. 3. <b>Warning:</b> Soil persistence of FARMALINX Metapris 500 SC Fungicide can be reduced under alkaline soil conditions. <b>Soil Spray:</b> a) Apply to soil surface immediately after sowing and repeat application at 10 weeks after sowing. b) Disease control will be improved if the seed is treated with FARMALINX Metapris 500 SC Fungicide prior to sowing. c) A further soil spray of 2 L/ha may be necessary if frequent or extended periods of cool, moist conditions occur later in the season. <b>Note:</b> 1. DO NOT spray directly over exposed seed in furrows before covering with soil. 2. <b>Warning:</b> Soil persistence of FARMALINX Metapris 500 SC Fungicide can be reduced under alkaline soil conditions. <b>Transplant Dip:</b> a) Dip seedlings for up to 4 hours in fungicide suspension before transplanting. b) A supplementary soil spray of 2 L/ha may be necessary if frequent or extended periods of cool, moist conditions occur in the late season.
		Qld, NSW, Vic, SA, NT only	2 L/ha in a minimum of 250 L of water		
		Qld, NSW, Vic, SA only	1 L/100 L of water		
Garlic	White Rot ( <i>Sclerotinia cepivorum</i> )	Qld, NSW, Vic, Tas, SA, NT only	10 mL/kg	-	<b>Pre-plant clove treatment:</b> Separate cloves, then add required amount of FARMALINX Metapris 500 SC Fungicide and mix thoroughly. <b>Warning:</b> Soil persistence of FARMALINX Metapris 500 SC Fungicide can be reduced under alkaline soil conditions.
Potato	Sclerotinia ( <i>Sclerotinia minor</i> )	All States	500 mL-1 L/ha	9 days	Apply at first spray just before hilling up. Apply a second spray just after hilling. Direct these sprays towards the stem bases and soil surface. Apply in sufficient water to ensure thorough coverage. Use the higher rate in situations where high disease levels are expected. Supplementary applications of 1 L/ha to foliage at 14-21 day intervals may be necessary if conditions favour further development of diseases.
	Target Spot ( <i>Alternaria solani</i> )		500 mL/ha		Apply in a program of sprays at 10 day intervals, beginning when warm weather conditions favour the disease and plants are 150-190 mm high. Apply in sufficient water to ensure thorough coverage.
Ornamentals	Sclerotinia Rot ( <i>Sclerotinia sclerotiorum</i> )	NSW only	75-100 mL/100 L water	-	Apply to run-off. Use the higher rate when disease is severe. DO NOT apply to African Violet flowers.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER THE APPROPRIATE LEGISLATION.**  
**DO NOT USE THIS PRODUCT IN THE HOME GARDEN.**

www.farmalinx.com

CONTINUED OVERLEAF

**RE-ENTRY PERIOD**  
*Re-entry* DO NOT enter treated area for 9 days after spray application unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing). Chemical resistant gloves and boots. Clothing must be laundered after each day's use.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**  
DO NOT contaminate streams, rivers or waterways with the chemical or used container or containers, which have held treated seed.

**STORAGE AND DISPOSAL**  
Store in a locked room or place away from children, animals, feedstuffs, seed and fertilisers. Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. The method of disposal of the container depends on the container type. Read the Storage and Disposal instructions on the label that is attached to the container.

**SAFETY DIRECTIONS**  
Contains procymidone which causes birth defects in laboratory animals. Women of child bearing age should avoid contact with procymidone. Very dangerous. Particularly if absorbed by skin contact, inhaled or swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. When opening the container, preparing product for use and using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and disposable mist mask or fume face mask covering mouth and nose. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

**FIRST AID**  
If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

**SDS**  
Additional information is listed in the safety data sheet (SDS). A safety data sheet for FARMALINX Metapris 500 SC Fungicide is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 9389 2455.

NOT re-use empty containers.  
DO NOT use treated seed for animal or human consumption. DO NOT allow treated seed to contaminate grain/other seed intended for animal or human consumption. DO NOT feed treated seed, or otherwise expose, to wild or domestic birds. Treated seed when stored should be kept apart from other grain and the containers should be clearly marked to indicate the contents have been treated with this product. DO NOT re-use empty containers.

**PRECAUTIONS**  
DO NOT use treated seed for animal or human consumption. DO NOT allow treated seed to contaminate grain/other seed intended for animal or human consumption. DO NOT feed treated seed, or otherwise expose, to wild or domestic birds. Treated seed when stored should be kept apart from other grain and the containers should be clearly marked to indicate the contents have been treated with this product. DO NOT re-use empty containers.

**COMPATIBILITY**  
This product is compatible with the following products: Agral, permethrin, propargite, pirimicarb and Avil. Provided the spray tank has an adequate agitation system and the mixture is not allowed to stand without agitation, these mixtures are acceptable. Some settling will occur in mixtures with permethrin and propargite.

**WETTING AGENT**  
Use a wetting agent at 10 mL/100 L of spray. With hard to wet crops the rate may be increased to 20 mL/100 L.

**Example Only**  
1. Dilute spray volume as determined above: For example 1500 L/ha  
2. Your chosen spray volume: For example 500 L/ha  
3. The concentration factor in this example is: 3X (i.e. 1500L divided by 500L = 3)  
4. If the dilute label rate is 10 mL/100 L, then the concentrate rate becomes 3 x 10, that is 30 mL/100 L of concentrate spray.  
The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows.  
• DO NOT use a concentrate rate higher than specified in the Critical Comment.  
• For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

**CONCENTRATE SPRAYING - Tree and Vine Crops only**  
• Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.  
• Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.  
• Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.  
• The mixing rate for concentrate spraying can then be calculated in the following way:  
Example Only  
1. Dilute spray volume as determined above: For example 1500 L/ha  
2. Your chosen spray volume: For example 500 L/ha  
3. The concentration factor in this example is: 3X (i.e. 1500L divided by 500L = 3)  
4. If the dilute label rate is 10 mL/100 L, then the concentrate rate becomes 3 x 10, that is 30 mL/100 L of concentrate spray.  
The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows.  
• DO NOT use a concentrate rate higher than specified in the Critical Comment.  
• For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off-label permits not endorsed by FARMALINX Pty Ltd or under abnormal conditions. FARMALINX Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

\* Other trademarks

**farmalinx**  
Partners in Agriculture

FARMALINX Pty Ltd ABN 95 134 353 245  
Level 25, Suite 2506, Tower 2,  
101 Grafton Street, Bondi Junction NSW 2022  
T: 02 9389 2455 F: 02 9389 2844  
Website: www.farmalinx.com

APVMA Approval No.: 69322/62976

**DANGEROUS POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**Metapris 500 SC**

FUNGICIDE  
ACTIVE CONSTITUENT: 500 g/L PROCYMIDONE

GROUP 2 FUNGICIDE

For the control of certain fungal diseases on various crops as per the Directions for Use table.

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT

FARMALINX Pty Ltd ABN 95 134 353 245  
Level 25, Suite 2506, Tower 2, 101 Grafton Street,  
Bondi Junction NSW 2022

**farmalinx**

**WITHHOLDING PERIODS:**  
**FABA BEANS, NAVY BEANS, STONE FRUIT (Blossom Bight control), WINE-GRAPES, POTATOES:**  
**DO NOT HARVEST FOR 9 DAYS AFTER APPLICATION.**  
**ONIONS:**  
**DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION.**

**GENERAL INSTRUCTIONS**  
**FUNGICIDE RESISTANCE WARNING**

FARMALINX Metapris 500 SC Fungicide is a member of the dicarboximide group of fungicides. For fungicide resistance management, this product is a Group 2 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 2 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product and other Group 2 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, FARMALINX Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

**MIXING**  
This product is suitable for application through conventional spray equipment calibrated to ensure thorough crop coverage. Add the required amount of product to the partially filled spray tank with the agitator running and complete filling the tank with water. Do not mix with alkaline water. Continue thorough agitation during spraying and after stoppage. DO NOT let prepared spray solution sit in the tank overnight.

**APPLICATION**  
**Dilute Spraying - Tree and Vine Crops only**  
• Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.  
• Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off. • The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.  
• Add the amount of product specified in the Directions for Use table for each 100 L of water. Spray to the point of run-off.  
• The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

GROUP 2 FUNGICIDE