



DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply by aerial application.
DO NOT treat vineyards that will or may be grazed by livestock.

CROP	DISEASE	RATE	WHP	CRITICAL COMMENTS
Grapevines	Bunch rot (<i>Botrytis cinerea</i>)	Dilute spray 120 g/100 L water Concentrate spray Refer to the Application section.	Harvest 4 weeks Grazing DO NOT treat vineyards that will or may be grazed by livestock	Apply as part of a <i>Botrytis</i> bunch rot program: In a 3 spray program, apply only 1 spray of RINKALS Fungicide. In a 4 spray program, apply only 2 sprays of RINKALS Fungicide. DO NOT apply consecutive sprays of RINKALS Fungicide. The ideal timing for the first spray is 5% capfall. DO NOT apply after 5% capfall on grapes grown for wine export. Applications can also be made at 80% capfall, pre-bunch closure, veraison or 4 weeks prior to harvest on table grapes or grapes for domestic wine. Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. DO NOT use in equipment that requires rates greater than 600 g/100 L (5X concentration). DO NOT apply in volumes less than 250 L/ha.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS:

GRAPEVINES: DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION.

GRAZING

GRAPEVINES: DO NOT TREAT VINEYARDS THAT WILL OR MAY BE GRAZED BY LIVESTOCK.

Export Slaughter Interval (ESI):

DO NOT FEED TREATED PRODUCE TO LIVESTOCK FOR 14 DAYS BEFORE SLAUGHTER

GENERAL INSTRUCTIONS

Farmalinx RINKALS Fungicide is a systemic anilide fungicide recommended for the control of Bunch rot (*Botrytis cinerea*) in grapevines. Optimum disease control is achieved when RINKALS Fungicide is applied in a regularly scheduled preventative spray program and is used in a rotational program with other effective fungicides. Because of its high specific activity, RINKALS Fungicide has good residual activity against target fungi.

EXPORT OF TREATED PRODUCE

Growers should note that suitable Maximum Residue Limits (MRLs) or import tolerances may not exist in all export markets for crops treated with RINKALS Fungicide. Additionally, some export markets have established MRLs different to those in Australia. If you are growing fruit for export (including fresh, dried or as wine) please check with your industry spray diary, peak industry body (such as the Australian Wine Research Institute www.awri.com.au) or FARMALINX for the latest information on MRLs and import tolerances before using RINKALS.

RESISTANT WEEDS WARNING

Farmalinx RINKALS Fungicide is a member of the oxathiin group of fungicides. For fungicide resistance management RINKALS is a Group 7 Fungicide. Some naturally occurring individual fungi resistant to the product and other Group 7 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. The resistant fungi will not be controlled by RINKALS Fungicide or other Group 7 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Farmalinx Pty Ltd accepts no liability for any losses that may result from the failure of RINKALS Fungicide to control resistant fungi.

GROUP 7 FUNGICIDE

MIXING

Partly fill the spray tank with water and add the required amount of product while stirring. If required, add compatible products and agitate thoroughly. Continue agitation as filling of the tank is completed.

Agitate again before spraying commences.

If tank mixing with other water dispersible granule (WG) or wettable powder (WP) formulations, ensure they are mixed in well before adding emulsifiable concentrate (EC) or suspension concentration (SC) products. WP formulations should be pre-mixed separately prior to adding to the spray tank.

APPLICATION

Dilute spraying

- Use a sprayer designed to apply high volumes of water up to the point of runoff and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of runoff. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes using different settings on the sprayer, from industry guidelines or expert advice.
- Add the amount of product specified in the Directions for Use table for each 100L of water. Spray to the point of run-off.
- The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate spraying

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of runoff) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.

The mixing rate for concentrate spraying can then be calculated in the following way:

EXAMPLE ONLY

Dilute spray volume as determined above: For example 1500 L/ha
Your chosen concentrate spray volume: For example 500 L/ha
The concentration factor in this example is: 3x (i.e. 1500 L ÷ 500L = 3)

If the dilute label rate is 120 g/100 L, then the concentrate rate becomes 3x 120, that is 360 g/100 L of concentrate spray.

- The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows.
- For concentrate application, DO NOT use in equipment that requires greater than 600 g/100 L (5x concentration). DO NOT apply in volumes less than 250L/ha.
- For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

COMPATIBILITY

RINKALS Fungicide is compatible with metiram (700 g/kg), mancozeb (750g/kg or 420g/L). RINKALS Fungicide is also compatible with a wide range of agricultural chemicals. A jar test with local water should be conducted before using any new tank mixtures with RINKALS Fungicide. For further information on compatibility please contact your local reseller or Farmalinx Pty Ltd representative.

RE-ENTRY PERIODS

DO NOT enter treated areas until the spray has dried unless wearing cotton overalls buttoned to the neck and the wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Toxic to aquatic life. DO NOT contaminate streams, rivers or watercourses with this product or used containers.

STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling,

break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. When opening the container and preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), a washable hat and elbow length PVC gloves. After each day's use, wash gloves and contaminated clothing. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MSDS

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FARMALINX RINKALS Fungicide is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 9389 2455.

NOTICE:

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Farmalinx Pty Ltd or under abnormal conditions. Farmalinx Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FARMALINX
Rinkals

FUNGICIDE
ACTIVE CONSTITUENT: 500 g/kg BOSCALID

GROUP 7 FUNGICIDE

For the control of bunch rot (*Botrytis cinerea*) in grapevines as specified in the Directions for Use

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.

FARMALINX Pty Ltd ABN 95 134 353 245
Level 25, Suite 2506, Tower 2, 101 Grafton Street,
Bondi Junction NSW 2022

farmalinx

farmalinx
Partners in Agriculture

FARMALINX Pty Ltd ABN 95 134 353 245

Level 25, Suite 2506, Tower 2,
101 Grafton Street, Bondi Junction NSW 2022

T: 02 9389 2455 F: 02 9389 2844

Website: www.farmalinx.com

APVMA Approval NO.: 81615/103463