

DIRECTIONS FOR USE**Restraints:**

DO NOT exceed 3 m between bait stations for mice or 9 m for rats. DO NOT place baits in open.

Place bait stations where mice and rats are active. Mice and rats require several bait stations. For effective control continue to bait until feeding stops.

Baiting Strategy:

Eliminate, as far as practicable, all alternative food sources. Bait infested area. Place bait stations under cover. Inspect bait stations daily. If eaten quickly increase number of bait stations. Continue observations and replenishment until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

Pellets are not subject to wind dispersal. Most deaths occur 4 to 7 days after consumption of bait. Locate bait stations along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Record the number and location of bait stations for purpose of replenishment and later removal.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

The bait is hazardous to pigs, cats, dogs, poultry, birds and wildlife and care should be exercised in its use. DO NOT place baits in locations which are accessible to domestic animals, livestock, birds or other animals. DO NOT contaminate dams, waterways and drains with bait or its used containers. The use of the product for the control of protected native rodents requires permission from wildlife authorities. The product is restricted to use in and around buildings (within 2 metres) or enclosed spaces e.g. drains.

STORAGE AND DISPOSAL

Containers which have held bait should not be used for any other purpose. Store in original container, tightly closed in a safe place out of reach of children. Dispose of empty containers and unused or untaken bait by burying in an approved dump. Look for dead mice and rats – burn or bury them, do not place in refuse bins or rubbish tips. Inspect all bait stations regularly and ensure baits are removed when rodent activity has ceased.

SAFETY DIRECTIONS

Poisonous if swallowed. DO NOT inhale dust. DO NOT touch bait, use scoop or measure. If on skin and after each baiting, wash thoroughly with soap and water. Vitamin K₁ (Phytomenadione) is antidotal.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. Vitamin K₁ (Phytomenadione) is antidotal.

NOTE TO PHYSICIANS AND VETERINARIANS:

Brodifacoum is a long lasting anticoagulant. Vitamin K₁ (Phytomenadione) should be administered urgently by medical or appropriate para-medical personnel. If administered intravenously the injection must be given slowly. The active constituent of Farmalinx RoDi Pellets is an anticoagulant chemical, which if accidentally taken by humans, domestic animals or pets, can reduce the clotting power of blood and haemorrhage may result.

MSDS

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FARMALINX RoDi Pellets is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 9389 2455.

NOTICE:

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Farmalinx Pty Ltd or under abnormal conditions. FARMALINX Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

farmalinx
Partners in Agriculture

FARMALINX Pty Ltd ABN 95 134 353 245

Level 25, Suite 2506, Tower 2,

101 Grafton Street, Bondi Junction NSW 2022

T: 02 9389 2455 F: 02 9389 2844

Website: www.farmalinx.com

APVMA Approval No.: 80059/100142

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FARMALINX

RoDi

PELLETS

ACTIVE CONSTITUENT: 0.05 g/kg BRODIFACOUm

For the control of rats and mice in and around industrial, agricultural and domestic buildings

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.

FARMALINX Pty Ltd ABN 95 134 353 245
Level 25, Suite 2506, Tower 2, 101 Grafton Street,
Bondi Junction NSW 2022

farmalinx