



READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FARMALINX

Sulfosulfuron 750 WG

HERBICIDE

ACTIVE CONSTITUENT: 750 g/kg SULFOSULFURON

GROUP B HERBICIDE

Selective herbicide for the control of certain weeds in wheat and triticale.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT

farmalinx *Partners in Agriculture*

FARMALINX Pty Ltd ABN 95 134 353 245
Level 25, Suite 2506, Tower 2, 101 Grafton Street, Bondi Junction NSW 2022
T: 02 9389 2455 F: 02 9389 2844 W: www.farmalinx.com

NET CONTENTS
500 g



DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply more than once per season

DO NOT apply to crops undersown with legumes

DO NOT spray when very dry conditions prevail

DO NOT use on furrow or flood-irrigated crops

DO NOT apply to soils with pH over 8.5

Weed control may be reduced if rainfall occurs soon after application

PRE-EMERGENT APPLICATION

CROP	WEEDS CONTROLLED/ SUPPRESSED	RATE	CRITICAL COMMENTS
Wheat Triticale (Vic, NSW, WA, SA, Tas only)	Barley grass* Brome grass* Wild turnip	25 g/ha	Apply to bare soil prior to sowing or at sowing and incorporate by the sowing operation. Application should not be made to ridged or excessively cloddy soil. In conservation tillage situations where weeds and grasses have emerged apply as a tank mixture with the recommended rate of 540 g/L Glyphosate (K salt) prior to sowing.
	Weeds as above, plus the control of Wild oats	25 g/ha plus 1.6 L/ha Tri-allate EC (500 g/L)	For best results follow the Tri-allate EC (500g/L) label directions for incorporation
	Weeds as above plus: Annual ryegrass Canary grass Fumitory Wireweed (Hogweed)	25 g/ha plus 800 mL/ha Trifluralin EC (480 g/L)	For best results follow the trifluralin label for incorporation

* *Suppression only*

POST-EMERGENT APPLICATION

Apply to wheat and triticale from early crop emergence to the 5 leaf/early tillering growth stage (Z11-15, 22)

USE THE SURFACTANT/WETTING AGENT – POST EMERGENT

Always add Bonza* at 1-2L/ha (1-2% volume/volume) of final spray volume

CROP	WEEDS CONTROLLED/ SUPPRESSED	STAGE OF WEED GROWTH	RATE	CRITICAL COMMENTS
Wheat Triticale (Vic, NSW, WA, SA, Tas only)	Amsinckia, Wild mustard (<i>Sinapsis arvensis</i>)*	Apply at the cotyledon to 4 leaf stage	25 g/ha	Optimum results will be obtained if good soil moisture exists at, and after, application. A follow-up spray with a different herbicide may be necessary to control subsequent germinations. When treating dense infestations of silver grass use water volumes of at least 70 L/ha and small droplets to improve coverage. FARMALINX SULFOSULFURON 750 WG Herbicide will provide good control of volunteer field peas. However a small proportion of plants may survive and require a follow-up spray with a different herbicide to eliminate the potential for grain contamination.
	Barley grass*, Silver grass (<i>Vulpia bromoides</i>), Wild oats*	Apply at the 1-4 leaf stage, (Z11-14)		
	Brome grass* (<i>Bromus diandrus</i> & <i>B. rigidus</i>)	Apply at the 1-3 leaf stage, (Z11-13)	For pH > 7.0 – 20 g/ha For pH < 7.0 – 25 g/ha	
	Volunteer canola (except canola tolerant to Group B herbicides), Wild radish, Wild turnip	Apply at the cotyledon to 4 leaf stage	20 g/ha	
	Volunteer field peas	Apply at the 1-3 leaf stage, (Z11-13)		

* Suppression only

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS
AUTHORISED UNDER APPROPRIATE LEGISLATION.**

WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.

CROP ROTATION GUIDELINES

Where the product is applied at the rate of 20 to 25 g/ha: Wheat and triticale can be planted the following season without restrictions. For other specified crops the FARMALINX SULFOSULFURON 750WG Herbicide treated area may be replanted after the interval indicated in the table below. These recommendations are made on the assumption that FARMALINX SULFOSULFURON 750WG Herbicide is applied to a wheat or triticale crop that reaches maturity in the season of application.

PLANT BACK PERIOD: Vic, NSW, WA, SA, TAS only

Soil pH (1:5 soil-water suspension method)	Replant interval	Minimum rainfall *	CROP
6.5 or less	10 months	300 mm	Canola, Chickpeas, Lentils, Lupins, Oats, Peas, Sub-clover **
	12 months	300 mm	Barley, Faba beans
6.5 to 8.5	10 months	300 mm	Cereal rye
	22 months	600 mm	Barley, Canola, Chickpeas, Faba beans, Lentils, Medic **, Oats, Peas, Vetch

* minimum rainfall required between FARMALINX SULFOSULFURON 750WG Herbicide application and the sowing of the plantback crop.

** Includes natural regeneration of sub-clover and medics. DO NOT apply to soils with a pH (water) > 8.5

For all other crops refer to FARMALINX for further advice.

GENERAL INSTRUCTIONS

FARMALINX SULFOSULFURON 750WG Herbicide is a selective herbicide absorbed by foliage and roots. It is rapidly translocated and acts by inhibiting acetolactate synthase (ALS), an enzyme necessary for root and shoot growth in sensitive plants. FARMALINX SULFOSULFURON 750WG Herbicide is a water dispersible granular herbicide for use in wheat and triticale. It can be used as a pre-plant, incorporated by sowing, for the control of wild turnip and suppression of barley grass and brome grass, or as a post-emergent for the control of silvergrass, wild radish and wild turnip, and suppression of barley grass, brome grass, wild mustard (*Sinapsis arvensis*) and wild oats.

PRE-EMERGENT

Apply to bare soil prior to sowing or at sowing and incorporate by the sowing operation to give thorough mixing. If applied to dry soil and sowing is to be delayed, incorporate immediately after spraying to prevent loss by wind erosion. For best results apply to moist soil when follow-up rain is likely to occur within 7-10 days. Weeds may emerge and will become stunted and uncompetitive soon after application although final results may not show for some weeks.

POST EMERGENT

Weeds should be young and actively growing. Use the higher rate under heavy weeds pressure and for larger weeds. A follow up spray with a suitable herbicide may be necessary to control subsequent germinations. Best weed control is obtained when rainfall wets the soil to a depth of 5 to 7.5cm within 7-10 days of application. Weeds will become stunted and uncompetitive soon after application although final results may not show for some weeks. When treatment is delayed or where weeds are not actively growing due to adverse conditions, for example, dry, waterlogged, frosty or diseased conditions, nutrient deficiency, high insect pressure or previous herbicide treatment, reduced levels of control may result. FARMALINX SULFOSULFURON 750WG Herbicide will remain in the soil for a period of time. The persistence of FARMALINX SULFOSULFURON 750WG Herbicide in the soil is dependent on various environmental conditions e.g. soil pH, soil moisture, soil temperature and organic

matter. Crops other than wheat and triticale may be very sensitive to low soil concentrations of FARMALINX SULFOSULFURON 750WG Herbicide, thus prior to using the product, careful consideration should be given to crop rotation plans (see Crop Safety and Crop Rotation Guidelines).

MIXING

FARMALINX SULFOSULFURON 750WG Herbicide is a water dispersible granular herbicide which mixes readily with water and is applied as a spray.

1. Partly fill the spray tank with water.
2. Start the agitation.
3. Add the correct amount of product to the spray tank with the agitation system running.
4. Continue agitation while topping up the tank with water and while spraying.
5. In tank mixes, FARMALINX SULFOSULFURON 750WG Herbicide must be in suspension before adding the companion product or surfactant (e.g. Bonza).
6. Use the spray mix within 24 hours of preparation.

APPLICATION

Apply by boom spray producing a MEDIUM spray quality. Ensure good spray coverage is obtained.

Apply using 40 to 100 litres of water per hectare. Avoid overlapping of boom runs. Use higher water volumes where crop or weed density is high.

DO NOT apply by aircraft.

SPRAYER CLEANUP

Where the sprayer is being used to spray wheat or triticale crops, rinse the sprayer thoroughly with water.

Where the sprayer is being used to spray crops other than wheat or triticale:

1. Drain tank and rinse tank and spray boom with clean water for at least 10 minutes.
2. Fill the tank with clean water and add to it 300mL of household chlorine bleach (containing 4% chlorine) per 100L of water. Rinse hoses and boom and leave in tank for 15 minutes whilst agitating. Drain through nozzle.
3. Repeat step 2 and then rinse thoroughly with clean water to remove all traces of chlorine bleach.
4. Nozzles and filters should be cleaned separately.
5. Dispose of all water used for cleaning.

Caution: DO NOT use chlorine bleach with ammonia. All traces of liquid fertiliser containing ammonia, ammonium nitrate or ammonium sulphate must be rinsed with water from the mixing and application equipment before adding chlorine bleach solution. Failure to do so will release a gas with a musty chlorine odour which can cause eye, nose and lung irritation. DO NOT clean equipment in an enclosed area.

COMPATIBILITY

Read and follow all label directions, restraints, plant-back and withholding periods and safety directions for all tank-mix products. In tank mixes, FARMALINX SULFOSULFURON 750WG Herbicide must be in suspension before adding the companion product or surfactant.

Pre-emergent:

FARMALINX SULFOSULFURON 750WG Herbicide is compatible with the following products:

- 500g/L Tri-allate,
- 450 g/L Glyphosate (IPA salt), 540 g/L Glyphosate (K salt), 680 g/kg Glyphosate (mono-ammonium salt), 250 g/L Paraquat dichloride,
- 135 g/L Paraquat dichloride/115 g/L Diquat dibromide,
- 480 g/L Trifluralin and Trifluralin in liquid hydrocarbon.

Post emergent:

FARMALINX SULFOSULFURON 750WG Herbicide is compatible with the following products:

- as a two way mixture with 300 g/L clopyralid (triisopropanolamine salt) (up to 300mL/ha)
- 500 g/L MCPA (isooctylester) (up to 350mL/ha)
- 250 g/L MCPA (ethylhexylester) + 25 g/L Diflufenican in 325 g/L liquid hydrocarbons/150 g/L n-methyl-2-pyrrolidone (up to 750mL/ha) for treatment of brome grass, wild oats, wild mustard (*Sinapsis arvensis*), wild radish and wild turnip. Some increased temporary crop yellowing may occur.

Always add Bonza at 1-2L/100L (1-2% volume/volume) of final spray volume.

FARMALINX SULFOSULFURON 750WG Herbicide may also be mixed with DC Trate at 2L/100L (2% volume/volume) of final spray volume. Some increased temporary crop yellowing may occur if FARMALINX SULFOSULFURON 750WG Herbicide is tank mixed with Nugrex/Tigrex.

Incompatible:

The following herbicides and insecticides are **incompatible** with FARMALINX SULFOSULFURON 750WG Herbicide:

400 g/kg Tralkoxydim WG

375 g/L and 500g/L Diclofop-methyl

500 g/L Terbutryn SC

290 g/L Omethoate

dicamba,

dimethoate

CROP SAFETY

DO NOT use on winter cereals undersown with legume pasture crops e.g. medics, clovers, lucerne.

DO NOT use on barley or oats. FARMALINX SULFOSULFURON 750WG Herbicide is effective on a range of plants, even at low rates of application. Since many rotational crops are sensitive, growers will need to consider the soil carry-over effects on subsequent crops. Below average rainfall, high levels of applied FARMALINX SULFOSULFURON 750WG Herbicide (greater than the recommended rates), pH greater than 6.5 and low soil temperatures and low rainfall all increase the carry-over risk. Where FARMALINX SULFOSULFURON 750WG Herbicide is used on soil types with pH greater than 8 (1:5 soil:water suspension method) and less than 300mm of rainfall has occurred since application, further advice should be sought from FARMALINX Pty Ltd regarding crop rotations, except for wheat and triticale.

Note: Some crop yellowing or crop retardation may occur where a stress factor such as water-logging, Rhizoctonia, take-all, cereal cyst nematode, nutrient deficiency or trace element deficiency is already present. Early season crop retardation may occur where the product is used on soils with a pH greater than 8 and which are prone to zinc deficiency.

See Crop Rotation Guidelines for further information.

RESISTANCE WEEDS WARNING

FARMALINX SULFOSULFURON 750WG Herbicide is a member of the sulfonylurea group of herbicides and has the ALS inhibitor mode of action. For weed resistance management, this product is a Group B herbicide. Some naturally-occurring weed biotypes resistant to FARMALINX SULFOSULFURON 750WG and other Group B herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by FARMALINX SULFOSULFURON 750WG or other Group B herbicides.

GROUP B HERBICIDE

To prevent or at least minimise the risk of resistant weeds occurring, use FARMALINX SULFOSULFURON 750WG tank mixes (if appropriate) and/or rotations with herbicides having different modes of action effective on the same weed species. DO NOT make more than one application of an ALS inhibitor herbicide to a crop, either pre-sowing incorporated by sowing or post crop and weed emergence. Since the occurrence of resistant weeds is difficult to detect prior to use, FARMALINX Pty Ltd accepts no liability for any losses that may result from the failure of FARMALINX SULFOSULFURON 750WG to control the resistant weeds. Advice as to strategies and alternative treatments that can be used should be obtained from your local farm chemical supplier, consultant, local Department of Agriculture, Department of Primary Industries, or a FARMALINX Pty Ltd representative.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply or drain or flush equipment on, or near desirable trees or other plants, where their roots may extend or in situations where by movement of soil, or seepage, absorption of the herbicide may occur. DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

Dangerous to plants, including aquatic plants.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate dams, streams, rivers or waterways with the chemical, or used containers.

STORAGE AND DISPOSAL

Store in closed, original containers in a cool, well-ventilated area away from fertilisers, pesticides and seeds. DO NOT store for prolonged periods in direct sunlight. DO NOT re-use containers. Triple rinse or (preferably) pressure rinse containers before disposal, and add rinsings to spray tank. DO NOT dispose of un-diluted chemical on-site. Break, crush or puncture the containers and bury at a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

SAFETY DIRECTIONS

Will irritate eyes. Avoid contact with eyes. DO NOT inhale dust. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre, phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FARMALINX SULFOSULFURON 750 WG Herbicide is available from Farmalinx Pty Ltd on request. Call Customer Service on 02 9389 2455.

NOTICE:

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Farmalinx Pty Ltd or under abnormal conditions. Farmalinx Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

*Registered Trade Mark

APVMA Approval No.: 69360/60589